



>>Variant: (±)-form

CAS Registry Number: 7685-44-1

Molecular Formula: C₅H₉NO₂

Molecular Weight: 115.132

Accurate Mass: 115.063329

Percentage Composition: C 52.16%; H 7.88%; N 12.17%; O 27.79%

Physical Description: Cryst. (MeOH)

Melting Point: Mp 243-245° (dec.)

Aldrich: 17344-4

Fluka: 5960

Sigma: A8378

>>Derivative: Me ester

CAS Registry Number: 70837-19-3

Molecular Formula: C₆H₁₁NO₂

Molecular Weight: 129.158

Accurate Mass: 129.078979

Percentage Composition: C 55.80%; H 8.58%; N 10.84%; O 24.77%

Physical Description: Liq.

Boiling Point: Bp_{1.1} 69-70°

>>Derivative: Et ester

CAS Registry Number: 89077-49-6

Extra CAS Registry Numbers(s): 68843-72-1

Molecular Formula: C₇H₁₃NO₂

Molecular Weight: 143.185

Accurate Mass: 143.094629

Percentage Composition: C 58.72%; H 9.15%; N 9.78%; O 22.35%

Physical Description: Liq.

Boiling Point: Bp₁₅ 75-78°

>>Derivative: *N*-Ac

CAS Registry Number: 50299-14-4

Molecular Formula: C₇H₁₁NO₃

Molecular Weight: 157.169

Percentage Composition: C 53.49%; H 7.05%; N 8.91%; O 30.54%

Melting Point: Mp 114°

Sigma: A7637

References:

- Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **1**, 574D, (*ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 874C; 875A, (*nmr*)
Black, S. *et al.*, *J. Biol. Chem.*, 1955, **213**, 39-50, (*synth, resoln*)
Japan. Pat., 1977, 77 54 085; *CA*, **87**, 150205, (*isol*)
Yaozhong, J. *et al.*, *Synth. Commun.*, 1989, **19**, 881-888, (*synth, ir, pmr*)
Sasaki, N.A. *et al.*, *Tet. Lett.*, 1989, **30**, 1943-1946, (*synth*)
Mooiweer, H.H. *et al.*, *Tetrahedron*, 1989, **45**, 4627-4636, (*synth, ir, pmr*)
Broxterman, Q.B. *et al.*, *J.O.C.*, 1992, **57**, 6286-6294, (*resoln, pmr, cmr*)
Dragovich, P.S. *et al.*, *J. Med. Chem.*, 1996, **39**, 1872-1884, (*Me ester, synth, ir, pmr*)
Org. Synth., 1999, **76**, 57-76, (*synth, ir, pmr, cmr*)