




>>Entry Name: 2-Amino-3-chloro-1,4-naphthoquinone

Synonym(s): 2-Amino-3-chloro-1,4-naphthalenedione, 9Cl. Quinoclamine, BSI. ACN, JMAF‡. ACNQ. Mogeton



CAS Registry Number: 2797-51-5

Molecular Formula: C₁₀H₆ClNO₂

Molecular Weight: 207.616

Accurate Mass: 207.008706

Percentage Composition: C 57.85%; H 2.91%; Cl 17.08%; N 6.75%; O 15.41%

Use/Importance: Has been used in herbicides

Physical Description: Yellow-brown needles (AcOH)

Melting Point: Mp 196-198° (193°)

Hazard & Toxicity: LD₅₀ (rat, orl) 1360 mg/kg

Rare Chemicals Library: S6576-6

>>Derivative: N-Ac

CAS Registry Number: 5397-78-4

Molecular Formula: C₁₂H₈ClNO₃

Molecular Weight: 249.653

Accurate Mass: 249.019271

Percentage Composition: C 57.73%; H 3.23%; Cl 14.20%; N 5.61%; O 19.23%

Physical Description: Red-brown platelets or yellow powder

Melting Point: Mp 219°

Rare Chemicals Library: S80256-5

>>Derivative: N-Ph

Synonym(s): 3-Anilino-2-chloro-1,4-naphthoquinone

CAS Registry Number: 1090-16-0

Molecular Formula: C₁₆H₁₀ClNO₂

Molecular Weight: 283.713

Accurate Mass: 283.040006

Percentage Composition: C 67.74%; H 3.55%; Cl 12.50%; N 4.94%; O 11.28%

Melting Point: Mp 214-216°

Aldrich: 24832-0

References:

- Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **2**, 81A, (*ir*)
Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 87A, (*nmr*)
Hoover, J.R.E. *et al.*, *J.A.C.S.*, 1954, **76**, 4148, (*synth*)
Gaultier, J. *et al.*, *Acta Cryst.*, 1965, **19**, 585, (*cryst struct*)
Neidlein, R. *et al.*, *Helv. Chim. Acta*, 1983, **66**, 2285, (*cmr*)
Hodnett, E.M. *et al.*, *J. Med. Chem.*, 1983, **26**, 570, (*synth*)
Hammam, A.S. *et al.*, *Coll. Czech. Chem. Comm.*, 1985, **50**, 71, (*synth, uv*)
Clark, N.G., *Pestic. Sci.*, 1985, **16**, 23, (*synth, activity*)
Pesticide Manual, 9th edn., 1991, No. 10517
Agrochemicals Handbook, 3rd edn., Royal Society of Chemistry, 1992, A1305
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AJ1250