



>>Variant: (±)-form

CAS Registry Number: 1115-47-5

Molecular Formula: C₇H₁₃NO₃S

Molecular Weight: 191.251

Accurate Mass: 191.061614

Percentage Composition: C 43.96%; H 6.85%; N 7.32%; O 25.10%; S 16.77%

Physical Description: Cryst. (H₂O)

Melting Point: Mp 114-115°

Aldrich: A1790-0

Sigma: A1626

>>Derivative: 4-(Acetylamino)phenyl ester

References:

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1280A, (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 784D, (ir)

Wheeler, G.P. *et al.*, *J.A.C.S.*, 1951, **73**, 4604, (synth)

U.S. Pat., 1962, 3 028 395; *CA*, **57**, 16742, (resoln)

Heyns, K. *et al.*, *Annalen*, 1963, **667**, 194, (ms)

U.K. Pat., 1967, 1 072 876; *CA*, **67**, 114015, (resoln)

Jung, G. *et al.*, *Eur. J. Biochem.*, 1973, **35**, 436, (cd)

Rotruck, J.T. *et al.*, *J. Nutr.*, 1975, **105**, 331, (metab)

Hawkes, G.E. *et al.*, *Nature (London)*, 1975, **257**, 767, (N-15 nmr)

Boggs, R.W., *Adv. Exp. Med. Biol.*, 1978, **105**, 571, (props)

Martindale, The Extra Pharmacopoeia, 28th/29th edn., Pharmaceutical Press, 1982

Ponnuswamy, M.N. *et al.*, *Acta Cryst. C*, 1985, **41**, 917, (cryst struct)

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ACQ275