



Aldrich: 85968-0

Fluka: 1010

Sigma: A6500

>>**Derivative:** Iodide

CAS Registry Number: 2260-50-6

Molecular Formula: $C_7H_{16}INO_2$

Molecular Weight: 273.113

Accurate Mass: 273.022577

Percentage Composition: C 30.78%; H 5.90%; I 46.47%; N 5.13%; O 11.72%

Melting Point: Mp 161°

Aldrich: 10043-9

Fluka: 1030

Sigma: A7000

References:

Aldrich Library of ^{13}C and 1H FT NMR Spectra, 1992, **1**, 1078A, (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 678C; 678D; 679A, (ir)

Dudley, H.W., *Biochem. J.*, 1929, **23**, 1064, (synth, bibl)

Marquardt, P. et al., *Arzneim.-Forsch.*, 1956, **6**, 168; 309, (biosynth)

Culvenor, C.C.J. et al., *Chem. Comm.*, 1966, 537, (pmr)

Beveridge, D.L. et al., *J.A.C.S.*, 1971, **93**, 3759, (struct)

Micelson, M.J. et al., *Acetylcholine; Approach of Mol. Mech. of Action*, Pergamon, 1974

Hanin, I., *Choline Acetylcholine: Handb. Chem. Assay Methods*, Raven Press, NY, 1974, (anal)

Svinning, T. et al., *Acta Cryst. B*, 1975, **31**, 1581, (cryst struct)

Sax, M. et al., *Acta Cryst. B*, 1976, **32**, 1953, (conformn)

Takasuka, M. et al., *J.C.S. Perkin 2*, 1982, 585, (ir)

Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 1112

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ABO000; CMF250