




>>Entry Name: Dimethylglyoxime

Synonym(s): 2,3-Butanedione dioxime. 2,3-Bis(hydroxyimino)butane. Biacetyl dioxime. Diethyl dioxime. Chugaev's reagent. DMG. H₂Dm



CAS Registry Number: 95-45-4

Related CAS Registry Numbers(s): 75006-64-3

Type of Compound Code(s): PA2400 PS8300 PA5010 PS7700 PM6850

Molecular Formula: C₄H₈N₂O₂

Molecular Weight: 116.119

Accurate Mass: 116.058578

Percentage Composition: C 41.37%; H 6.94%; N 24.12%; O 27.56%

Use/Importance: Used in detn. of Ni, Pd; gravimetric detn. of Ni (red pptn.), Pd (yellow), photometric detn. of Ni (λ_{max} 445 nm, ϵ 15000), Pd, Co, Fe, Cu, Re; in separation of Ni, Pd by extraction (CHCl₃) or pptn. as M(HDm)₂

Physical Description: Triclinic cryst. (EtOH aq.)

Melting Point: Mp 238-240°

Solubility: Sol. EtOH, Me₂CO, Et₂O, alk. soln.; v. spar. sol. H₂O, CHCl₃

Aldrich: 16257-4

Fluka: 40390

Sigma: D2783

>>Derivative: O,O'-Di-Ac

CAS Registry Number: 19800-34-1

Molecular Formula: C₈H₁₂N₂O₄

Molecular Weight: 200.194

Accurate Mass: 200.079708

Percentage Composition: C 48.00%; H 6.04%; N 13.99%; O 31.97%

Physical Description: Cryst. (EtOH)

Melting Point: Mp 112.5-113.5°

>>Derivative: O,O'-Dibenzoyl

CAS Registry Number: 18128-12-6

Molecular Formula: C₁₈H₁₆N₂O₄

Molecular Weight: 324.335

Accurate Mass: 324.111008

Percentage Composition: C 66.66%; H 4.97%; N 8.64%; O 19.73%

Physical Description: Needles

Melting Point: Mp 225°

References:

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 489C, (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 922D, (ir)

Aldrich Library of NMR Spectra, **10**, 136A, (pmr)

Chugaev, L.A., *Ber.*, 1905, **38**, 2520, (synth, props)

Org. Synth., Coll. Vol., 2, 1943, 204, (synth)

Christopherson, H. et al., *Anal. Chim. Acta*, 1954, **10**, 1, (detn, Ni)

Norwitz, G. et al., *Anal. Chem.*, 1965, **37**, 417, (use, Ni)

Talati, A.M. et al., *J. Indian Chem. Soc.*, 1968, **45**, 242, (uv)

Davis, W.F., *Talanta*, 1969, **16**, 1330, (use, Pd)

Schienbaum, M.L., *J.O.C.*, 1970, **35**, 2790, (synth)

Keresztury, G. et al., *Spectrochim. Acta A*, 1977, **33**, 29, (ir, Raman)

Singh, R.B. et al., *Talanta*, 1979, **26**, 425, (rev)

Raston, C.L. et al., *Aust. J. Chem.*, 1980, **33**, 1519, (cryst struct)

Zav'yalov, S.I. et al., *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1984, 161; *Bull. Acad. Sci. USSR, Div. Chem. Sci. (Engl. Transl.)*, 1984, 142, (synth)

Kinoshita, S. et al., *Bull. Chem. Soc. Jpn.*, 1986, **59**, 653, (cmr)

Marczenko, Z., *Separation and Spectrophotometric Determination of Elements*, Ellis Horwood, Chichester, 1986, 393

Onishi, H., *Photometric Determination of Traces of Metals, Part IIb: Individual Metals. Magnesium to Zinc*, John Wiley, New York, 1989, 125

Schildcrout, S.M. et al., *Inorg. Chem.*, 1990, **29**, 1054, (ms)

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DBH000