



>>Entry Name: Disodium phosphorofluoridate, 9Cl, 8Cl

Synonym(s): Sodium monofluorophosphate(V). Disodium fluorotrioxophosphate(V)

CAS Registry Number: 10163-15-2

Extra CAS Registry Numbers(s): 7631-97-2

$\text{Na}_2[\text{P}(\text{O})_3\text{F}]$

Molecular Formula: $\text{FNa}_2\text{O}_3\text{P}$

Molecular Weight: 143.95

Accurate Mass: 143.936444

Percentage Composition: F 13.20%; Na 31.94%; O 33.34%; P 21.52%

Source/Synthesis: Synth. from $\text{HF} + \text{NaPO}_3$, or $\text{Na}_4\text{P}_2\text{O}_7$; or from $\text{NaF} + \text{NaPO}_3$ or $\text{Na}_3\text{P}_3\text{O}_9$; or from $\text{Ag}_2\text{PO}_3\text{F} + \text{NaCl}$

Use/Importance: Caries inhibiting component in toothpaste

Melting Point: Mp 625°

Solubility: Sol. H_2O (4 g per 100 cm^3). Insol. EtOH, Et_2O

Other Data: In H_2O at ca. 400° $\rightarrow$ HF + diphosphate

Aldrich: 34444-3

Fluka: 71528

>>Derivative: Decahydrate

Source/Synthesis: Formed from aq. solns. at 0-5°

>>Variant: α -form

Molecular Formula: $\text{FNa}_2\text{O}_3\text{P}$

Molecular Weight: 143.95

Accurate Mass: 143.936444

Percentage Composition: F 13.20%; Na 31.94%; O 33.34%; P 21.52%

General Statement: Orthorhombic

Source/Synthesis: Synth. by dehydration of hydrate

>>Variant: β -form

Molecular Formula: $\text{FNa}_2\text{O}_3\text{P}$

Molecular Weight: 143.95

Accurate Mass: 143.936444

Percentage Composition: F 13.20%; Na 31.94%; O 33.34%; P 21.52%

General Statement: Orthorhombic. The Na coordination polyhedron is a distorted octahedron. Struct. contains two types of PO_3F tetrahedra

Source/Synthesis: Synth. by cryst. from aq. soln. at <20°

References:

Inorg. Synth., 1950, **3**, 106-109, (synth)

Gmelin Handbook Inorg. Chem., Syst. No. 21, 1967, **4**, 1675, (rev, bibl)

Singh, E.B. *et al.*, *J. Indian Chem. Soc.*, 1970, **47**, 491, (uv)

U.K. Pat., 1970, 1 186 664; *CA*, **72**, 134776, (manuf)

Reddy, G.S. *et al.*, *Z. Naturforsch., B*, 1970, **25**, 1199, (F-19, P-31 nmr)

Durand, J. *et al.*, *Acta Cryst. B*, 1974, **30**, 1565, (synth, cryst struct)

Fluck, E. *et al.*, *Pure Appl. Chem.*, 1975, **44**, 373, (pe)

Granier, W. *et al.*, *Rev. Chim. Miner.*, 1975, **12**, 147, (synth, props, cryst struct)

Martynchik, S.P. *et al.*, *Izv. Akad. Nauk SSSR, Neorg. Mater.*, 1981, **17**, 703; *Inorg. Mater. (Engl. Transl.)*, 1981, **17**, 474, (synth, ir, hydrol)

Kirillov, S.A. *et al.*, *Zh. Neorg. Khim.*, 1982, **27**, 25; *Russ. J. Inorg. Chem. (Engl. Transl.)*, 1982, **27**, 14, (ir, Raman)

White, W.E., *Caries Res., Suppl.*, 1983, **17**, 2, (rev)

Haubenreisser, U. *et al.*, *Mol. Phys.*, 1987, **60**, 151, (P-31 nmr)

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DXD600