




>>Entry Name: Dopastin

Synonym(s): *N*-[2-(Hydroxynitrosoamino)-3-methylbutyl]-2-butenamide, 9CI



Type of Compound Code(s): XA1950 XA3950 XA1650

Molecular Formula: C₉H₁₇N₃O₃

Molecular Weight: 215.252

Accurate Mass: 215.126992

Percentage Composition: C 50.22%; H 7.96%; N 19.52%; O 22.30%

Use/Importance: Dopamine β-hydroxylase inhibitor. Antihypertensive, antineoplastic agent

Solubility: BERDY SOL: Sol. MeOH, C₆H₆; fairly sol. hexane; poorly sol. H₂O

UV: [acid] λ_{max} 215 (ε 20900) (0.01*N* HCl) (Derep) [base] λ_{max} 213 (ε 17800); 246 (ε 10800) (0.01*N* NaOH) (Derep) [neutral] λ_{max} 213 (ε 18100); 245 (ε 10500) (pH 7 phosphate buffer) (Derep) [neutral] λ_{max} 213 (E1%/1cm 840); 245 (E1%/1cm 190) (pH 7 buffer) (Berdy) [acid] λ_{max} 215 (E1%/1cm 970) (HCl) (Berdy) [base] λ_{max} 213 (E1%/1cm 330); 246 (E1%/1cm 500) (NaOH) (Berdy)

>>Variant: (*S,E*)-form

CAS Registry Number: 37134-80-8

Type of Compound Code(s): VV0130 AH0130

Molecular Formula: C₉H₁₇N₃O₃

Molecular Weight: 215.252

Accurate Mass: 215.126992

Percentage Composition: C 50.22%; H 7.96%; N 19.52%; O 22.30%

Biological Source: Obt. from *Pseudomonas* bacteria

Melting Point: Mp 116-119°

Optical Rotation: [α]_D²¹ -246 (c, 0.5 in EtOH)

pKa Value: p*K*_a 5.1

Other Data: Pharmacol. active isomer

Hazard & Toxicity: LD₅₀ (mus, orl) 750 mg/kg. LD₅₀ (mus, ipr) 460 mg/kg

References:

Ger. Pat., 1972, *Microbiochemical Research Foundation*, 2 214 757; *CA*, **77**, 163039y, (*pharmacol*)

Ohno, M. *et al.*, *Chem. Comm.*, 1973, 147, (*abs config*)

Umezawa, H., *Pure Appl. Chem.*, 1973, **33**, 129, (*pharmacol*)

Iinuma, H. *et al.*, *Agric. Biol. Chem.*, 1974, **38**, 2099, (*synth, props*)