



**Percentage Composition:** C 49.28%; H 4.87%; F 27.51%; N 6.76%; O 11.58%

>>**Derivative:** Acetate salt

**Synonym(s):** Flecainide acetate, USAN. Flecaine. Modicard. Tambocor. R 818

**CAS Registry Number:** 54143-56-5

**Physical Description:** Granular solid (diisopropyl ether)

**Melting Point:** Mp 145-147°

**Hazard & Toxicity:** LD<sub>50</sub> (rat, orl) 1346 mg/kg

**Sigma:** F6777

**References:**

- Banitt, E.H. *et al.*, *J. Med. Chem.*, 1977, **20**, 821; 1986, **29**, 299, (*synth, pharmacol, resoln, bibl*)  
Huduk, J.M. *et al.*, *Am. J. Cardiol.*, 1984, **53**, 17B, (*rev*)  
Blaschke, G. *et al.*, *Chem. Ber.*, 1985, **118**, (*resoln, abs config*)  
Holmes, B. *et al.*, *Drugs*, 1985, **29**, 1, (*rev*)  
Broly, F. *et al.*, *Biomed. Chromatogr.*, 1991, **5**, 248, (*hplc*)  
Alessi-Severini, S. *et al.*, *Clin. Chem. (Winston-Salem, N.C.)*, 1991, **37**, 111, (*hplc, enantiomers*)  
Alessi-Severini, S. *et al.*, *Anal. Profiles Drug Subst.*, 1992, **21**, 169, (*rev*)  
Fischer, C. *et al.*, *Ther. Drug Monit.*, 1992, **14**, 433, (*gc-ms, hplc*)  
Banitt, E.H. *et al.*, *Chron. Drug Discovery*, 1993, **3**, 163, (*rev*)  
Yamashita, T. *et al.*, *J. Pharmacol. Exp. Ther.*, 1995, **274**, 315, (*pharmacol*)  
Zipes, D.P., *Am. J. Cardiol.*, 1996, **71**, 1A; 53A, (*use, rev*)  
Heisler, B.E. *et al.*, *J. Pharmacol. Exp. Ther.*, 1996, **279**, 317, (*pharmacol*)  
*Martindale, The Extra Pharmacopoeia*, 31st edn., Pharmaceutical Press, 1996, 868