



Accurate Mass: 191.055798
Percentage Composition: C 50.27%; H 4.22%; F 29.82%; N 7.33%; O 8.37%
Boiling Point: Bp₁₀ 57-60°
Refractive Index: n_D²⁰ 1.4520
Aldrich: 34098-7

>>Derivative: *O,N*-Di-Me

CAS Registry Number: 702-24-9
Molecular Formula: C₉H₁₃NO
Molecular Weight: 151.208
Accurate Mass: 151.099714
Percentage Composition: C 71.49%; H 8.67%; N 9.26%; O 10.58%
Boiling Point: Bp₁₄ 121°

>>Derivative: *O,N,N*-Tri-Me

CAS Registry Number: 15175-54-9
Molecular Formula: C₁₀H₁₅NO
Molecular Weight: 165.235
Accurate Mass: 165.115364
Percentage Composition: C 72.69%; H 9.15%; N 8.48%; O 9.68%
Boiling Point: Bp₁₆ 110-111°

>>Derivative: *O,N,N*-Tri-Me; hydrochloride

CAS Registry Number: 61185-92-0
Melting Point: Mp 157°

References:

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 1288D, (*ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 610A, (*nmr*)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1172D, (*ir*)
Salkowski, H., *Ber.*, 1889, **22**, 2137
Tiffeneau, M., *Bull. Soc. Chim. Fr.*, 1911, **9**, 819
Ingold, C.K. *et al.*, *J.C.S.*, 1929, 1199, (*deriv*)
Witkop, B. *et al.*, *J.A.C.S.*, 1954, **76**, 5589
Larsen, P.O., *Biochim. Biophys. Acta*, 1965, **107**, 134, (*isol, uv*)
Olsen, O. *et al.*, *Phytochemistry*, 1980, **19**, 1783, (*isol, pmr, cmr*)
Larsen, P.O. *et al.*, *Phytochemistry*, 1984, **23**, 895, (*isol*)
Eur. Pat., 1990, Mitsui, 354 297; *CA*, **113**, 59234j
Beutler, J.A. *et al.*, *J. Antibiot.*, 1990, **43**, 107, (*BMV 40660*)
Thomas, S. *et al.*, *J.O.C.*, 2001, **66**, 1999-2004, (*Me ether, pmr, cmr*)