Molecular Formula: $C_{28}H_{34}O_9$
Molecular Weight: 514.571
Accurate Mass: 514.220285
Percentage Composition: C 65.36%; H 6.66%; O 27.98%
Biological Source: Isol. from *Schizandra chinensis* and *Schizandra sphenanthera*
Use/Importance: Antihepatitic agent. Shows insecticidal props.
Physical Description: Cryst.
Melting Point: Mp 95-97°
Optical Rotation: $[\alpha]_D^{23}$ –27.6 (c, 0.5 in CH_2Cl_2)
UV: [neutral] λ_{max} 218 (ε 94500); 257 (sh) (ε 17400); 292 (sh) (ε 4470) (EtOH) (Derep)

>>**Derivative:** 7,8-Diepimer, 7-benzoyl
Synonym(s): Gomisin C. Schizantherin A. Schisantherin A. Wuweizi ester A

CAS Registry Number: 58546-56-8

Molecular Formula: $C_{30}H_{32}O_9$
Molecular Weight: 536.577
Accurate Mass: 536.204635
Percentage Composition: C 67.15%; H 6.01%; O 26.84%
Biological Source: Isol. from *Schizandra chinensis* and *Schizandra sphenanthera*
UV: [neutral] λ_{max} 220 (ε 57500); 253 (sh) (ε 15500); 284 (sh) (ε 3800) (EtOH) (Derep)

References:

Taguchi, K. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 3296, *(isol)*
Liu, C.-S. *et al.*, *Sci. Sin. (Engl. edn.)*, 1978, **21**, 483, *(isol)*
Ikeya, Y. *et al.*, *Chem. Pharm. Bull.*, 1979, **27**, 1383; 1576; 1980, **28**, 3357; 1990, **38**, 1408, *(isol, ms, cmr, struct)*
Hikino, H. *et al.*, *Planta Med.*, 1984, **50**, 213, *(pharmacol)*
Miyazawa, M. *et al.*, *Nat. Prod. Lett.*, 1998, **12**, 175-180, *(Gomisin B, pmr, cmr)*