




>>Entry Name: *N*-Leucyltyrosine



Molecular Formula: C₁₅H₂₂N₂O₄

Molecular Weight: 294.35

Accurate Mass: 294.157958

Percentage Composition: C 61.21%; H 7.53%; N 9.52%; O 21.74%

>>**Variant:** (*S,S*)-form

Synonym(s): L-L-form

CAS Registry Number: 968-21-8

Molecular Formula: C₁₅H₂₂N₂O₄

Molecular Weight: 294.35

Accurate Mass: 294.157958

Percentage Composition: C 61.21%; H 7.53%; N 9.52%; O 21.74%

Physical Description: Cryst. + 2H₂O

Melting Point: Mp 275-277°

Optical Rotation: [α]_D²⁰ +11.3

Solubility: Sol. H₂O, spar. sol. EtOH

pKa Value: pK_{a1} 3.46; pK_{a2} 7.84; pK_{a3} 10.09 (25°)

Sigma: L0501

>>**Derivative:** *N*-*tert*-Butyloxycarbonyl, Me ester

Melting Point: Mp 113-114°

>>**Derivative:** *N*-Benzyloxycarbonyl-*O*-*tert*-butyl methyl ester

Melting Point: Mp 87-89°

>>**Variant:** (*R*)-(*S*)-form

Synonym(s): D-Leucyl-L-tyrosine

CAS Registry Number: 3303-29-5

Molecular Formula: C₁₅H₂₂N₂O₄

Molecular Weight: 294.35

Accurate Mass: 294.157958

Percentage Composition: C 61.21%; H 7.53%; N 9.52%; O 21.74%

Physical Description: Powder + 2H₂O (Et₂O/EtOH)

Melting Point: Mp 250-260 dec.°

Optical Rotation: [α]_D²⁰ -16

pKa Value: pK_{a3} 10.35

Sigma: L3877

References:

Abderhalden, E. *et al.*, *Chem. Zentralbl.*, 1930, **2**, 2271; 1931, **1**, 2210

Kinoshita, M. *et al.*, *Annalen*, 1966, **696**, 226

Pettit, G.R. *et al.*, *Org. Mass Spectrom.*, 1970, **3**, 67