



>>Entry Name: Lithium perchlorate (LiClO₄)

Synonym(s): Lithium chlorate(VII). Chloric(VII) acid, lithium salt. Perchloric acid, lithium salt, 8Cl

CAS Registry Number: 7791-03-9

LiClO₄

Molecular Formula: ClLiO₄

Molecular Weight: 106.391

Accurate Mass: 105.964515

Percentage Composition: Cl 33.32%; Li 6.52%; O 60.15%

Source/Synthesis: Prepd. by slow neutralisation of Li₂CO₃ with aq. HClO₄; cryst. as trihydrate

Use/Importance: Commercially available. Component of rocket propellants

Physical Description: Colourless orthorhombic cryst., deliquescent. Dec. on heating with evolution of O₂ and formation of LiCl

Melting Point: Mp 236°

Solubility: V. sol. H₂O (60 g per 100 cm³ at 25°; 150 g per 100 cm³ at 89°), EtOH (152 g per 100 cm³ at 25°), MeOH (182 g per 100 cm³ at 25°), Me₂CO (137 g per 100 cm³ at 25°), Et₂O (114 g per 100 cm³ at 25°)

Other Data: ΔH_f° -381 kJ mol⁻¹; ΔG_f° -254 kJ mol⁻¹; S° 126 J K⁻¹ mol⁻¹

Hazard & Toxicity: Oxidiser, irritant; potentially explosive when mixed with org. compds.

Aldrich: 20528-1

Fluka: 62585

Sigma: L7656

>>Derivative: Monohydrate

CAS Registry Number: 18716-61-5

Molecular Formula: ClH₂LiO₅

Molecular Weight: 124.407

Accurate Mass: 123.97508

Percentage Composition: Cl 28.50%; H 1.62%; Li 5.58%; O 64.30%

Other Data: ΔH_f° -697 kJ mol⁻¹; ΔG_f° -510 kJ mol⁻¹; S° 155 J K⁻¹ mol⁻¹

>>Derivative: Trihydrate

Synonym(s): Perchloric acid, lithium salt trihydrate, 8Cl

CAS Registry Number: 13453-78-6

Molecular Formula: ClH₆LiO₇

Molecular Weight: 160.437

Accurate Mass: 159.99621

Percentage Composition: Cl 22.10%; H 3.77%; Li 4.33%; O 69.81%

Source/Synthesis: Prep. by electrolysis of aq. LiClO₃ soln.

Use/Importance: Commercially available. Pyroelectric, piezoelectric

Physical Description: White hexagonal cryst.

Melting Point: Mp 95°

Solubility: V. sol. H₂O (130 g per 100 cm³ at 25°), EtOH (73 g per 100 cm³ at 25°), MeOH (156 g per 100 cm³ at 25°), Me₂CO (96 g per 100 cm³ at 25°); prac. insol. Et₂O (0.096 g per 100 cm³ at 25°)

Other Data: ΔH_f° -1298 kJ mol⁻¹; ΔG_f° -1001 kJ mol⁻¹; S° 255 J K⁻¹ mol⁻¹

Hazard & Toxicity: Oxidiser. Dec. on heating by H₂O loss prior to explosive reaction

Aldrich: 20530-3

Fluka: 62589

Sigma: L3533

References:

- Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **2**, 1253B, (*ir*)
West, C.D., *Z. Kristallogr.*, 1934, **88**, 198, (*cryst struct, trihydrate*)
Gluyas, R.E., *Diss. Abstr.*, 1957, **17**, 2800, (*cryst struct*)
Gmelin Handbook Inorg. Chem., Syst. No. 20, 1960, 404, (*rev, bibl*)
Hezel, A. *et al.*, *Spectrochim. Acta*, 1966, **22**, 1949, (*ir*)
Sequeira, A. *et al.*, *Acta Cryst., Sect. B*, 1975, **31**, 1735, (*cryst struct, trihydrate*)
Berglund, B. *et al.*, *J. Mol. Struct.*, 1978, **43**, 169, (*ir, trihydrate*)
Lundgren, J.O. *et al.*, *Acta Cryst., Sect. B*, 1982, **38**, 15, (*cryst struct, trihydrate*)
Chase, M.W. *et al.*, *J. Phys. Chem. Ref. Data*, 1985, **14**, 763, (*thermochem data*)
Emmeneger, F. *et al.*, *Inorg. Chem.*, 1997, **36**, 3146-3151, (*soln*)
Sigma-Aldrich Library of Chemical Safety Data, **2**, 2156D; 2157A
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, LHN000