



>>Entry Name: Mercury(II) fluoride (HgF₂)

Synonym(s): Mercuric fluoride

CAS Registry Number: 7783-39-3

HgF₂

Molecular Formula: F₂Hg

Molecular Weight: 238.587

Accurate Mass: 239.967423

Percentage Composition: F 15.93%; Hg 84.07%

Source/Synthesis: Synth. from HgCl₂ and F₂ at 100°, or from Hg(OAc)₂ and SF₄, or by heating Hg₂F₂ at 450° or from Hg₂F₂ and Cl₂ or Br₂

Use/Importance: Fluorinating agent; used in optical fibres

Physical Description: Light sensitive colourless cryst.

Melting Point: Mp 645 dec.°

Boiling Point: Bp 650°

Solubility: Sol. HF, dil. HNO₃. Dec. in H₂O

Aldrich: 33932-6

Fluka: 83386

>>Derivative: Dihydrate

CAS Registry Number: 26453-89-4

Source/Synthesis: Synth. from red HgO and HF

Physical Description: Cryst.

Other Data: Struct. contains 8-coord. Hg (4F + 4O) in a Thompson cube. Readily hydrol. forming FHgOH

References:

- Ebert, F. *et al.*, *Z. Anorg. Allg. Chem.*, 1933, **210**, 269
Swanson, H.E. *et al.*, *NBS Monogr. (U.S.)*, 1963, **25**, Sect. 2, (*powder struct*)
Gmelin Handbook Inorg. Chem., Syst. No. 34, 1967, **B2**, 408, (*rev, bibl*)
Loewenschuss, A. *et al.*, *J. Chem. Phys.*, 1969, **50**, 2502, (*ir, vapour*)
Polishchuk, S.A. *et al.*, *J. Less-Common Met.*, 1970, **21**, 63, (*synth, ir, pmr, thermal anal, dihydrate*)
Gagarinskii, Yu.V. *et al.*, *Spectrosc. Lett.*, 1970, **3**, 23, (*F-19 nmr, salt and dihydrate*)
Vaughan, R.W. *et al.*, *J. Chem. Phys.*, 1972, **57**, 5383, (*F-19 nmr*)
Bukvetskii, B.V. *et al.*, *Koord. Khim.*, 1976, **2**, 1208, (*cryst struct, dihydrate*)
Kirk-Othmer Encycl. Chem. Technol., 3rd edn., Wiley, 1978, **10**, 763
Poole, R.T. *et al.*, *Phys. Rev. B: Condens. Matter*, 1979, **20**, 1733, (*pe*)
Givan, A. *et al.*, *J. Chem. Phys.*, 1980, **72**, 3809, (*ir, Raman, matrix*)
Shendrik, V.P. *et al.*, *Ukr. Khim. Zh. (Russ. edn.)*, 1982, **48**, 1108, (*synth*)
Laubach, S. *et al.*, *Hyperfine Interact.*, 1985, **23**, 259, (*Hg-199 Mössbauer*)