




>>Entry Name: 2-Oxazolidinone, 9Cl, 8Cl



CAS Registry Number: 497-25-6

Molecular Formula: C₃H₅NO₂

Molecular Weight: 87.078

Accurate Mass: 87.032029

Percentage Composition: C 41.38%; H 5.79%; N 16.09%; O 36.75%

Melting Point: Mp 89°

Aldrich: O-940-9

Fluka: 75793

>>Derivative: N-Ac

CAS Registry Number: 1432-43-5

Molecular Formula: C₅H₇NO₃

Molecular Weight: 129.115

Accurate Mass: 129.042594

Percentage Composition: C 46.51%; H 5.46%; N 10.85%; O 37.17%

Physical Description: Cryst. (C₆H₆/Et₂O)

Melting Point: Mp 69-70°

Aldrich: 34851-1

>>Derivative: N-Me

CAS Registry Number: 19836-78-3

Molecular Formula: C₄H₇NO₂

Molecular Weight: 101.105

Accurate Mass: 101.047679

Percentage Composition: C 47.52%; H 6.98%; N 13.85%; O 31.65%

Boiling Point: Bp₁ 120°

Hazard & Toxicity: LD₅₀ (rat, orl) approx. 7 g/kg

Aldrich: 22800-1

Rare Chemicals Library: S53693-8

References:

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 774A; 774B, (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1261C; 1262A; 1262C, (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 787C, (*ir*)

Dyen, M.E. *et al.*, *Chem. Rev.*, 1967, **67**, 197, (*rev*)

Simons, S.S. *et al.*, *J.O.C.*, 1973, **38**, 414, (*synth*)

Sonoda, N. *et al.*, *Tet. Lett.*, 1975, 1969, (*synth*)

Hammer, C.F. *et al.*, *J.O.C.*, 1981, **46**, 1521, (*synth, ir, pmr*)

Lown, J.W. *et al.*, *J.O.C.*, 1983, **48**, 3901, (*deriv, synth, ir, pmr, cmr, ms*)

Shibata, I. *et al.*, *Bull. Chem. Soc. Jpn.*, 1989, **62**, 853, (*synth, derivs*)

Wouters, J. *et al.*, *Acta Cryst. C*, 1997, **53**, 895-897, (*cryst struct*)

Wouters, J. *et al.*, *Acta Cryst. C*, 1997, **53**, 895-897, (*cryst struct*)

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MND750; OMM000