




>>Entry Name: Pentolium(2+)

Synonym(s): 1,1'-(1,5-Pentanediy)bis-1-methylpyrrolidinium, 9Cl. 1,1-Pentamethylenebis-1-methylpyrrolidinium, 8Cl. Pentapyrrolidine. Pentolinium. Pentolon. Pentolonium. Pyrroplegium. Many other names



CAS Registry Number: 144-44-5

Molecular Formula: $C_{15}H_{32}N_2^{2(+)}$

Molecular Weight: 240.431

Accurate Mass: 240.256548

Percentage Composition: C 74.93%; H 13.41%; N 11.65%

Use/Importance: Antihypertensive, ganglion blocking agent

Other Data: Used mainly as hydrogen tartrate

>>Derivative: Hydrogen tartrate

Synonym(s): Pentolinium tartrate, BAN. Pentolonium tartrate, INN. Ansolysen

CAS Registry Number: 52-62-0

Molecular Formula: $C_{23}H_{42}N_2O_{12}$

Molecular Weight: 538.591

Accurate Mass: 538.273778

Percentage Composition: C 51.29%; H 7.86%; N 5.20%; O 35.65%

Melting Point: Mp 203 dec.°

Solubility: Sol. H_2O , insol. Et_2O , $CHCl_3$

Hazard & Toxicity: LD₅₀ (rat, orl) 890 mg/kg; LD₅₀ (rat, ivn) 50.8 mg/kg

Aldrich: 86146-4

Sigma: P3520

>>Derivative: Diiodide

Molecular Formula: $C_{15}H_{32}I_2N_2$

Molecular Weight: 494.24

Accurate Mass: 494.065494

Percentage Composition: C 36.45%; H 6.53%; I 51.35%; N 5.67%

Physical Description: Cryst. + $\frac{1}{2} H_2O$

Melting Point: Mp 292°

References:

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 396B, (ir)

Libman, D.D. *et al.*, *J.C.S.*, 1952, 2305, (synth)

Sammul, O.R. *et al.*, *J. Assoc. Off. Agric. Chem.*, 1964, **47**, 918, (ir)

Barafsky, D.F. *et al.*, *Int. J. Mass Spectrom. Ion Phys.*, 1983, **46**, 359, (ms)

Negwer, M., *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 3796, (synonyms)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 382

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PBT000