



>>Entry Name: Phosphorus chloride (PCl₅)

Synonym(s): Phosphorus pentachloride. Pentachlorophosphorane, 9Cl. Phosphoric chloride. Tetrachlorophosphorus(1+) chloride

CAS Registry Number: 10026-13-8

PCl₅ (vap.) PCl₄⁽⁺⁾PCl₆⁽⁻⁾ or PCl₄⁽⁺⁾Cl⁽⁻⁾ (solid state)

Molecular Formula: Cl₅P

Molecular Weight: 208.237

Accurate Mass: 205.818022

Percentage Composition: Cl 85.13%; P 14.87%

Use/Importance: Catalyst in org. reactions. Chlorinating agent, e.g. converting many primary and secondary alcohols to alkyl chlorides; amides → chloroimines.

Deoxygenating agent e.g. for nitro compds. Commercially available

Hazard & Toxicity: Corrosive. Attacks eyes and mucous membranes

Aldrich: 15777-5

Fluka: 79590

>>Variant: Molecular-form

Molecular Formula: Cl₅P

Molecular Weight: 208.237

Accurate Mass: 205.818022

Percentage Composition: Cl 85.13%; P 14.87%

polar solvs. in equilibrium with ionic forms

Other Data: Forms a weak complex with C₆H₆. In vap., some dissociation → PCl₃ + Cl₂

>>Variant: Ionic-form

Synonym(s): Tetrachlorophosphonium hexachlorophosphate. Tetrachlorophosphorus(1+) hexachlorophosphate(1-)

CAS Registry Number: 19453-01-1

Molecular Formula: Cl₁₀P₂

Molecular Weight: 416.474

Accurate Mass: 411.636044

Percentage Composition: Cl 85.13%; P 14.87%

polar solvs. e.g. MeCN, MeNO₂, PhNO₂ and also CCl₄

Physical Description: Moisture-sensitive off white or pale yellowish cryst. which fume strongly evolving HCl gas. Pungent odour

Melting Point: Mp 167°

Boiling Point: Subl. 160°

Other Data: At lower concs. (<0.03M) in polar solvs., the struct. reverts to [PCl₄⁽⁺⁾][Cl⁽⁻⁾], which is also the struct. in HCl(l)

References:

Aldrich Library of Infrared Spectra, 3rd edn., 1981, 554E, (ir)

Gmelin Handbook Inorg. Chem., Syst. No. 16, 1965, 434, (rev, bibl)

DiLorenzo, J.V. *et al.*, *Inorg. Chem.*, 1967, **6**, 766, (nqr)

Mellor Compr. Treat. Inorg. Theor. Chem., 1971, **8/III**, 464; 476; 1147, (rev, bibl)

Preiss, H., *Z. Anorg. Allg. Chem.*, 1971, **380**, 51; 1972, **389**, 280, (cryst struct, ms)

Takahashi, Y., *Bull. Chem. Soc. Jpn.*, 1972, **45**, 4, (pe)

Fichter, M., *Spectrochim. Acta B*, 1975, **30**, 417, (pe)

Fridland, S.V. *et al.*, *Usp. Khim.*, 1978, **47**, 1397; *Russ. Chem. Rev. (Engl. Transl.)*, 1978, **47**, 742, (rev, bibl)

Cahay, R. *et al.*, *J. Chem. Phys.*, 1979, **70**, 5534; 5539, (phase diagram, Raman, bibl)

Ripmeester, J.A. *et al.*, *J. Chem. Phys.*, 1979, **70**, 5545; 1980, **73**, 1474, (P-31 nmr, Raman)

Sergienko, L.M. *et al.*, *Zh. Obshch. Khim.*, 1979, **49**, 1982; *J. Gen. Chem. USSR (Engl. Transl.)*, 1979, **49**, 1743, (uv)

McLlolland, B.W. *et al.*, *J. Mol. Struct.*, 1983, **99**, 309, (ed)

Finch, A. *et al.*, *Polyhedron*, 1986, **5**, 1537, (Raman)

Bay, E. *et al.*, *J.O.C.*, 1988, **53**, 2858, (use)

Harris, R.K. *et al.*, *Mol. Phys.*, 1989, **66**, 993, (P-31 nmr)

Finch, A. *et al.*, *Chem. Comm.*, 1993, 957, (metastable form, cryst struct)

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PHR500