



UV: [neutral] $\lambda_{\max}$ (EtOH) (Berdy)

Sigma: A8905

>>Derivative: 15-Ac, 4-O- α -D-glucopyranoside

CAS Registry Number: 99124-45-5

Molecular Formula: C₂₃H₃₄O₁₁

Molecular Weight: 486.515

Accurate Mass: 486.210115

Percentage Composition: C 56.78%; H 7.04%; O 36.17%

Biological Source: From *Fusarium sulphureum*

Physical Description: Pale-yellow gum

Optical Rotation: $[\alpha]_{\text{D}}^{20} +79.5$ (c, 0.585 in MeOH)

Other Data: Less toxic than other trichothecenes

>>Derivative: 4,15-Di-Ac

Synonym(s): Diacetoxyscirpenol. Anguidin. ANG 66. NSC 141537

CAS Registry Number: 2270-40-8

Molecular Formula: C₁₉H₂₆O₇

Molecular Weight: 366.41

Accurate Mass: 366.167855

Percentage Composition: C 62.28%; H 7.15%; O 30.57%

Biological Source: Constit. of *Fusarium* spp.

Physical Description: Cryst. (EtOAc)

Melting Point: Mp 161-162°

Optical Rotation: $[\alpha]_{\text{D}}^{19} +20$ (c, 1 in Me₂CO)

Calculated Log P Data: Log P -0.12 (uncertain value) (calc)

UV: [neutral] $\lambda_{\max}$ 0 (end) (ϵ) (MeOH) (Derep)

Hazard & Toxicity: Very toxic if swallowed. Skin irritant. LD₅₀ (rat, orl) 7 mg/kg. Exp. reprod. and teratogenic effects. Exp. immunotoxicological props. Other exp. effects. Associated with outbreaks of mycotoxicoses in farm animals

Sigma: D0761

>>Derivative: Tri-Ac

Molecular Formula: C₂₁H₂₈O₈

Molecular Weight: 408.447

Accurate Mass: 408.17842

Percentage Composition: C 61.75%; H 6.91%; O 31.34%

Physical Description: Cryst. (EtOH)

Melting Point: Mp 124-125°

Optical Rotation: $[\alpha]_{\text{D}}^{23} +44$ (c, 1 in Me₂CO)

References:

- Sigg, H.P. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 962, (struct)
Tidd, B.K., *J.C.S.(C)*, 1967, 218, (pmr)
Bamburg, J.R. *et al.*, *Tetrahedron*, 1968, **24**, 3329, (deriv)
Cole, M. *et al.*, *Appl. Microbiol.*, 1972, **24**, 660, (use)
Ripperger, H. *et al.*, *Phytochemistry*, 1975, **14**, 2298, (isol)
Pathre, S.V. *et al.*, *J. Agric. Food Chem.*, 1976, **24**, 97, (isol)
Ellison, R.A. *et al.*, *J.O.C.*, 1976, **41**, 576, (cmr)
Steyn, P.S. *et al.*, *Phytochemistry*, 1978, **17**, 949, (isol)
Hartmann, G.R. *et al.*, *Planta Med.*, 1978, **34**, 231, (rev, Anguidin)
Claridge, C.A. *et al.*, *Appl. Environ. Microbiol.*, 1979, **37**, 693-696, (3-Ac)
Brumley, W.C. *et al.*, *Biomed. Mass Spectrom.*, 1982, **9**, 451, (ms)
Pohland, A.E. *et al.*, *Pure Appl. Chem.*, 1982, **54**, 2220, (uv, ir, pmr, ms, cd)
Brooks, D.W. *et al.*, *J.A.C.S.*, 1983, **105**, 4472, (synth, deriv)
Trichothecenes - Chemical, Biological and Toxicological Aspects, (ed. Ueno, Y.), Elsevier, 1983, (tox, rev)
Gorst-Allman, C.P. *et al.*, *J.C.S. Perkin 1*, 1985, 1553, (deriv)
Roush, W.R. *et al.*, *J.O.C.*, 1985, **50**, 3224, (deriv)
Anderson, D.W. *et al.*, *Tet. Lett.*, 1987, **28**, 2661, (Anguidin)
Ziegler, F.E. *et al.*, *J.A.C.S.*, 1990, **112**, 2749, (synth)
Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, New York, 1981, 160; 168; 171
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AOP250