




>>Entry Name: Stigmasta-7,22-dien-3-one, 9Cl
ENTRY UNDER REVIEW



Molecular Formula: C₂₉H₄₆O
Molecular Weight: 410.682
Accurate Mass: 410.354865
Percentage Composition: C 84.81%; H 11.29%; O 3.90%

>>Variant: (5 α ,22*E*,24*S*)-form
Synonym(s): α -Spinasterone

CAS Registry Number: 23455-44-9

Molecular Formula: C₂₉H₄₆O
Molecular Weight: 410.682
Accurate Mass: 410.354865
Percentage Composition: C 84.81%; H 11.29%; O 3.90%
Biological Source: Isol. from heartwood of *Albizzia julibrissin* and bark of *Acacia concinna*
Physical Description: Needles (Me₂CO)
Melting Point: Mp 170-172° (164-166°)
Optical Rotation: [α]_D²⁰ +23

>>Derivative: Oxime

Molecular Formula: C₂₉H₄₇NO
Molecular Weight: 425.696
Accurate Mass: 425.365764
Percentage Composition: C 81.82%; H 11.13%; N 3.29%; O 3.76%
Physical Description: Cryst. (Me₂CO)
Melting Point: Mp 239-241°

References:

Breton, J.L. *et al.*, *J.C.S.*, 1963, 1401, (*synth*)
Mahato, S.B. *et al.*, *Phytochemistry*, 1972, **11**, 2116, (*isol*, *synth*)
Nakano, Y. *et al.*, *Mokuzai Gakkaishi*, 1975, **21**, 577; *CA*, **84**, 102290w, (*isol*)
Iyer, C. *et al.*, *Phytochemistry*, 1978, **17**, 2036, (*isol*)
Brooks, C.J.W. *et al.*, *Lipids*, 1980, **15**, 745, (*synth*)
Herath, H.M.T.B. *et al.*, *ACGC Chem. Res. Commun.*, 1999, **9**, 3-8, (*isol*)
Akihisa, T. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1161-1163, (*pmr*, *cmr*)