



CAS Registry Number: 2010-15-3
Molecular Formula: C₉H₈N₂OS
Molecular Weight: 192.241
Accurate Mass: 192.035733
Percentage Composition: C 56.23%; H 4.19%; N 14.57%; O 8.32%; S 16.68%
Physical Description: Yellow plates (EtOH)
Melting Point: Mp 240-242 dec.°
Hazard & Toxicity: LD₅₀ (mus, orl) 155 mg/kg

Rare Chemicals Library: S67654-3
Sigma: P0627

>>Derivative: 1,3-*N*-Di-Me

CAS Registry Number: 1801-62-3
Molecular Formula: C₅H₈N₂OS
Molecular Weight: 144.197
Accurate Mass: 144.035733
Percentage Composition: C 41.65%; H 5.59%; N 19.43%; O 11.10%; S 22.24%
Physical Description: Cryst. (C₆H₆/petrol)
Solubility: V. sol. hot H₂O, sol. EtOH, Et₂O, CHCl₃, C₆H₆

References:

- Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **1**, 1348B, (*nmr*)
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 836B, (*ir*)
Johnson, T.B. *et al.*, *J.A.C.S.*, 1915, **37**, 2406, (*synth*)
Dalglish, C.E. *et al.*, *J.C.S.*, 1947, 559, (*synth*)
Edward, J.T., *Org. Sulfur Compd.*, 1966, **2**, 287, (*rev*)
Walker, L.A., *Acta Cryst. B*, 1969, **25**, 88, (*cryst struct*)
Rowley, G.L. *et al.*, *J.A.C.S.*, 1971, **93**, 5542, (*deriv, synth*)
Thielman, H., *Z. Chem.*, 1978, **18**, 174, (*synth*)
Blotny, G., *Synthesis*, 1983, 391, (*deriv*)
Reddick, R.E. *et al.*, *J.A.C.S.*, 1987, **109**, 4380, (*deriv, synth*)
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, ADC750; TFJ750