




>>Entry Name: Thonzylamine, BAN, INN

Synonym(s): *N*-[(4-Methoxyphenyl)methyl]-*N*',*N*'-dimethyl-*N*-2-pyrimidinyl-1,2-ethanediamine, 9Cl. 2-[(2-Dimethylaminoethyl)-(*p*-methoxybenzyl)amino]pyrimidine, 8Cl



CAS Registry Number: 91-85-0

Molecular Formula: C₁₆H₂₂N₄O

Molecular Weight: 286.376

Accurate Mass: 286.179361

Percentage Composition: C 67.11%; H 7.74%; N 19.56%; O 5.59%

Use/Importance: Potent antihistamine

Development Status: Discontinued

Boiling Point: Bp₆ 202-204°. Bp_{2.2} 185-187°

Hazard & Toxicity: LD₅₀ (mus, orl) 245 mg/kg. Exp. reprod. effects

>>Derivative: Hydrochloride

Synonym(s): Thonzylamine hydrochloride, INN. Anohist. Neohetramine. Resistab. Many other names

CAS Registry Number: 63-56-9

Melting Point: Mp 173-176°

pKa Value: pK_{a1} 8.84; pK_{a2} 2.08 (25°)

Hazard & Toxicity: LD₅₀ (mus, orl) 245 mg/kg

Sigma: T2771

>>Derivative: Hydrobromide

Melting Point: Mp 159-160.5°

>>Derivative: Dipicrate

Physical Description: Yellow needles

Melting Point: Mp 141-145°

References:

U.S. Pat., 1949, 2 465 865; *CA*, **43**, 6244

Saijo, S., *Yakugaku Zasshi*, 1952, **72**, 1444; *CA*, **47**, 8080

Iwasaki, T., *Nippon Yakurigaku Zasshi*, 1957, **53**, 375; *CA*, **52**, 14850e, (*pharmacol*)

Shell, J.W. *et al.*, *Mikrochim. Acta*, 1960, 31

Chang, C.-J. *et al.*, *J. Pharm. Sci.*, 1976, **65**, 1019, (*cmr*)

Mossini, F. *et al.*, *Farmaco, Ed. Sci.*, 1984, **39**, 189, (*synth, activity*)

Negwer, M., *Organic-Chemical Drugs and their Synonyms*, 6th edn., *Akademie-Verlag*, 1987, 4096

Parvez, M., *Acta Cryst. C*, 1988, **44**, 1433

Martindale, The Extra Pharmacopoeia, 30th edn., *Pharmaceutical Press*, 1993, 946

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., *Van Nostrand Reinhold*, 1992, NCD500; RDU000