



>>Entry Name: Tributylborane, 10Cl, 9Cl

CAS Registry Number: 122-56-5

$\text{B}(\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3)_3$

Molecular Formula: $\text{C}_{12}\text{H}_{27}\text{B}$

Molecular Weight: 182.156

Accurate Mass: 182.22058

Percentage Composition: C 79.13%; H 14.94%; B 5.94%

Use/Importance: Additive used in polym. of alkenes. Reagent for transfer of a butyl group. Commercially available

Physical Description: Liq.

Boiling Point: Bp_{20} 108-110°. Bp_9 90-91°

Hazard & Toxicity: May ignite spontaneously esp. as thin film or if absorbed on paper. LD_{50} (rat, orl) 1125 mg/kg

Aldrich: 17990-6

>>Derivative: Me_3N complex

Synonym(s): Tributyl(trimethylamine)boron. Tributyl(*N,N*-dimethylmethanamine)boron, 10Cl. *N,N*-Dimethylmethanamine compd. with tributylborane (1:1)

CAS Registry Number: 62685-92-1

Molecular Formula: $\text{C}_{15}\text{H}_{36}\text{BN}$

Molecular Weight: 241.267

Accurate Mass: 241.294079

Percentage Composition: C 74.67%; H 15.04%; B 4.48%; N 5.81%

References:

Johnson, J.R. *et al.*, *J.A.C.S.*, 1938, **60**, 115, (*synth*)

Costes, J.P. *et al.*, *J. Chim. Phys. Phys.-Chim. Biol.*, 1976, **73**, 16, (*deriv*)

Compr. Org. Chem., (Eds. Barton, D.H.R. *et al*), Pergamon, Oxford, 1979, **3**, 792, (*rev*)

Kuimova, M.E. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1979, 314, (*nmr*)

Brown, H.C. *et al.*, *J. Organomet. Chem.*, 1980, **188**, 1, (*use*)

Akira, S., *Acc. Chem. Res.*, 1982, **15**, 178, (*rev, uses*)

Akira, S., *Acta Cryst.*, 1982, **15**, 178, (*rev, uses*)

Brown, H.C. *et al.*, *Tetrahedron*, 1987, **43**, 4059, (*reactions*)

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, THX500

Bretherick, L., *Handbook of Reactive Chemical Hazards*, 4th edn., Butterworths, 1990, 3321