>>**Entry Name: 1,2,3,4-Tetrahydro-6,7-dimethoxyisoquinoline,** 9Cl
Synonym(s): Heliamine



CAS Registry Number: 1745-07-9

Molecular Formula: $C_{11}H_{15}NO_2$
Molecular Weight: 193.245
Accurate Mass: 193.110279
Percentage Composition: C 68.37%; H 7.82%; N 7.25%; O 16.56%
Biological Source: Alkaloid from *Pachycereus weberi, Pachycereus pringlei, Pachycereus pecten-aboriginum, Backebergia militaris* and *Carnegiea gigantea* (Cactaceae)
Biological Use/Importance: Antineoplastic agent. Inhibitor of rat sarcoma 45
Melting Point: Mp 84-85°
Calculated Log P Data: Log P 0.98 (calc)
Other Data: Readily forms carbonate on exp. to air

>>**Derivative:** Hydrochloride

CAS Registry Number: 2328-12-3
Physical Description: Cryst. (EtOH/Et_2O)
Melting Point: Mp 255° (248°, 252°)
Aldrich: 29191-9

>>**Derivative:** Oxalate

Melting Point: Mp 214-215°

>>**Derivative:** Picrate

Physical Description: Cryst. (EtOH)
Melting Point: Mp 223-225°

>>**Derivative:** *N*-Me

References:

Buck, J.S. *et al.*, *J.A.C.S.*, 1934, **56**, 1769-1771, *(synth)*
Bobbitt, J.M. *et al.*, *J.O.C.*, 1965, **30**, 2247-2250, *(synth)*
Hughes, D.W. *et al.*, *Can. J. Chem.*, 1976, **54**, 2252-2260, *(cmr)*
Strömbom, J. *et al.*, *Acta Pharm. Suec.*, 1978, **15**, 127-132; *CA*, **89**, 126107w, *(isol)*
Mata, R. *et al.*, *Phytochemistry*, 1980, **19**, 673-678; 1983, **22**, 1263-1270, *(isol, ir, uv, pmr, cmr, ms, di-Me ether)*
Pummangura, S. *et al.*, *J. Nat. Prod.*, 1982, **45**, 277-282, *(isol)*
Ruchirawat, S. *et al.*, *Synth. Commun.*, 1984, **14**, 1221-1228, *(synth, pmr, ms)*
Murahashi, S.I. *et al.*, *Bull. Chem. Soc. Jpn.*, 1989, **62**, 2968, *(pmr, cmr)*
Shinohara, T. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 813-819, *(synth)*

https://www.baizuoyan.com - CopyRight ©baizuoyan