



>>Entry Name: Sulfuryl chloride, 11Cl
Synonym(s): Sulfur(VI) dichloride dioxide

CAS Registry Number: 7791-25-5
 $\text{S(O)}_2\text{Cl}_2$

Molecular Formula: $\text{Cl}_2\text{O}_2\text{S}$
Molecular Weight: 134.97
Accurate Mass: 133.899604
Percentage Composition: Cl 52.53%; O 23.71%; S 23.76%

Source/Synthesis: Synth from $\text{SO}_2 + \text{Cl}_2$
Use/Importance: Commercially available. Used as a chlorinating agent
Physical Description: Colourless liq., hydrol. to HCl and H_2SO_4
Melting Point: Mp -54°
Boiling Point: Bp 69.2°
Solubility: Sol. C_6H_6 , CHCl_3 , Et_2O
Density: d_4^0 1.705
Refractive Index: n_D^{20} 1.4430
Hazard & Toxicity: Corrosive to skin, toxic, lachrymator. Strong odour

Aldrich: 32052-8
Fluka: 86211

References:

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 1227A, (ir)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1692B, (ir)
Inorg. Synth., 1939, **1**, 114, (synth)
Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1967, **1**, 1128; 1969, **2**, 394; 1972, **3**, 276; 1974, **4**, 474; 1975, **5**, 641; 1977, **6**, 561; 1982, **10**, 375, (uses)
Brauer, G., *Handbuch Prap. Anorg. Chem.*, 3rd edn., Ferdinand Enke Verlag, 1975, **1**, 388, (synth)
Kirk-Othmer Encycl. Chem. Technol., 3rd edn., Wiley, 1978, **22**, 131
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, SOT000