




>>Entry Name: 1,2,3,4-Tetrahydro-2-methyl-4-phenyl-8-isoquinolinamine, 9CI

Synonym(s): 8-Amino-1,2,3,4-tetrahydro-2-methyl-4-phenylisoquinoline. **Nomifensine**, BAN



CAS Registry Number: 24526-64-5

Molecular Formula: C₁₆H₁₈N₂

Molecular Weight: 238.332

Accurate Mass: 238.146998

Percentage Composition: C 80.63%; H 7.61%; N 11.75%

Use/Importance: Thymoleptic, CNS stimulant

Development Status: No longer marketed (withdrawn due to toxicity)

Calculated Log P Data: Log P 2.66 (calc)

Sigma: N1530

>>Variant: (±)-form

Molecular Formula: C₁₆H₁₈N₂

Molecular Weight: 238.332

Accurate Mass: 238.146998

Percentage Composition: C 80.63%; H 7.61%; N 11.75%

Melting Point: Mp 179-181°

>>Derivative: Maleate

Synonym(s): **Nomifensine maleate**, USAN. Alival. Many other names

CAS Registry Number: 32795-47-4

Physical Description: Cryst. (EtOH)

Melting Point: Mp 199-201°

Hazard & Toxicity: Adverse effects incl. risk of haemolytic anaemia and renal failure. LD₅₀ (rat, orl) 430 mg/kg. Exp. reprod. effects

References:

U.K. Pat., 1969, 1 164 192; CA, **72**, 21621e, (synth)

Maj, J. et al., *Arzneim.-Forsch.*, 1976, **26**, 1109, (pharmacol)

Eiden, F. et al., *Dtsch. Apoth. -Ztg.*, 1977, **117**, 611, (props, spectra)

Fields, E.D., *Drug Intell. Clin. Pharm.*, 1982, **16**, 547, (pharmacol)

Linford Rees, W., *Int. Congr. Symp. Ser. R. Soc. Med.*, R. Soc. Med., London, 1984, **70**, (book)

Heptner, W. et al., *J. Clin. Psychiatry*, 1984, **45**, 12; 21; 33, (rev)

Berova, N. et al., CA, 1985, **103**, 141804x, (synth, resoln, abs config)

Zava-Kaczian, E. et al., *J. Med. Chem.*, 1986, **29**, 1189, (synth)

Negwer, M., *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 3981

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 266

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, NMV700; NMV725