




>>Entry Name: Xenytropium(1+)

Synonym(s): 8-([1,1'-Biphenyl]-4-ylmethyl)-3-(3-hydroxy-1-oxo-2-phenylpropoxy)-8-methyl-8-azoniabicyclo[3.2.1]octane, 9Cl. 3-Hydroxy-8-(*p*-phenylbenzyl)tropanium, 8Cl. 8-(*p*-Phenylbenzyl)atropinium. Fentropilium. Dendrepar. Gastripon. Gastropin. BTTB. FX 501. N 399



Related CAS Registry Numbers(s): 112967-95-0 112967-96-1 112967-98-3

Molecular Formula: C₃₀H₃₄NO₃⁽⁺⁾

Molecular Weight: 456.603

Accurate Mass: 456.253869

Percentage Composition: C 78.92%; H 7.51%; N 3.07%; O 10.51%

>>Variant: (±)-form

Molecular Formula: C₃₀H₃₄NO₃

Molecular Weight: 456.603

Accurate Mass: 456.253869

Percentage Composition: C 78.92%; H 7.51%; N 3.07%; O 10.51%

>>Derivative: Chloride

CAS Registry Number: 112967-97-2

Physical Description: Cryst. (MeOH/EtOH)

Melting Point: Mp 236-238°

>>Derivative: Bromide

Synonym(s): Xenytropium bromide, INN

CAS Registry Number: 511-55-7

Molecular Formula: C₃₀H₃₄BrNO₃

Molecular Weight: 536.507

Accurate Mass: 535.192205

Percentage Composition: C 67.16%; H 6.39%; Br 14.89%; N 2.61%; O 8.95%

Use/Importance: Antimuscarinic, spasmolytic agent

Physical Description: Cryst. (MeOH)

Melting Point: Mp 230° (224-225°)

Hazard & Toxicity: LD₅₀ (rat, orl) 1880 mg/kg. LD₅₀ (rat, ipr) 66 mg/kg

References:

Gyermek, L. *et al.*, *Acta Physiol. Acad. Sci. Hung.*, 1955, **8**, 43, (*pharmacol*)

U.S. Pat., 1958, 2 833 773; *CA*, **52**, 15602b, (*synth, pharmacol*)

Suga, T. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 2235, (*metab*)

Nador, K. *et al.*, *Arzneim.-Forsch.*, 1987, **37**, 1324, (*synth, pmr, derivs*)

Martindale, The Extra Pharmacopoeia, 30th edn., *Pharmaceutical Press*, 1993, 433

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., *Van Nostrand Reinhold*, 1992, PEM750