



>>Entry Name: Undecanal

ENTRY UNDER REVIEW

Synonym(s): Undecyl aldehyde. Undecylic aldehyde. FEMA 3092

CAS Registry Number: 112-44-7

$\text{H}_3\text{C}(\text{CH}_2)_9\text{CHO}$

Molecular Formula: $\text{C}_{11}\text{H}_{22}\text{O}$

Molecular Weight: 170.294

Accurate Mass: 170.167065

Percentage Composition: C 77.58%; H 13.02%; O 9.40%

Biological Source: Found in many essential oils, e.g. *Citrus* spp., *Cunila lythrifolia* and kumquat *Fortunella marginata*

Use/Importance: Perfumery and flavouring ingredient

Melting Point: Mp -4°

Boiling Point: Bp₁₈ 116-117°

Density: d_4^{23} 0.825

Refractive Index: n_D^{23} 1.4322

Other Data: Readily polymerises

Hazard & Toxicity: Skin irritant. LD₅₀ (rat, orl) >5000 mg/kg

Aldrich: W30920-6

Fluka: 94020

Sigma: U8252

Supelco: S48-3092

>>Derivative: Oxime

Molecular Formula: $\text{C}_{11}\text{H}_{23}\text{NO}$

Molecular Weight: 185.309

Accurate Mass: 185.177964

Percentage Composition: C 71.30%; H 12.51%; N 7.56%; O 8.63%

Physical Description: Cryst. (MeOH)

Melting Point: Mp 72°

>>Derivative: Semicarbazone

Physical Description: Cryst. (MeOH)

Melting Point: Mp 103°

>>Derivative: 2,4-Dinitrophenylhydrazone

Physical Description: Yellow cryst. (EtOH)

Melting Point: Mp 104°

References:

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 732C, (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 469B, (ir)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 555D, (ir)

Blaise, E.E., *Bull. Soc. Chim. Fr.*, 1904, **31**, 492, (synth)

Allen, C.F.H. *et al.*, *J.O.C.*, 1941, **6**, 596, (synth)

Davies, R.R. *et al.*, *J. Soc. Chem. Ind., London*, 1943, **62**, 128, (synth)

Org. Synth., 1971, **51**, 39, (synth)

Opdyke, D.L.J., *Food Cosmet. Toxicol.*, 1973, **11**, 481, (rev, tox)

Lewis, R.J., *Food Additives Handbook*, Van Nostrand Reinhold International, New York, 1989, UJJ000

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992