



>>Entry Name: Antimony chloride oxide (SbClO), 9Cl, 8Cl

Synonym(s): Chlorooxostibine, 10Cl. Chlorooxoantimony(III)

CAS Registry Number: 7791-08-4

SbClO

Molecular Formula: ClOSb

Molecular Weight: 173.202

Accurate Mass: 171.867588

Percentage Composition: Cl 20.47%; O 9.24%; Sb 70.29%

General Statement: Monoclinic. Consists of puckered sheets in which 2/3 of the Sb atoms are linked to 2 oxygens and 1 chlorine, and the remaining Sb atoms to 4 oxygens, the $(\text{Sb}_6\text{O}_6\text{Cl}_4)^{2(+)}$ sheets being held together by $\text{Cl}^{(-)}$ ions. Two types of Cl atoms distinguished by Sb-Cl length, 230-250 pm within the polymeric layer and 290 pm between Sb and inter-layer Cl

Source/Synthesis: Synth. from $\text{SbCl}_3 + \text{H}_2\text{O}$

Use/Importance: Fireproofing agent

Physical Description: Colourless cryst. or cryst. powder

Solubility: Sol. aq. HCl and aq. tartaric acid

Other Data: Dec. in $\text{H}_2\text{O} \rightarrow \text{Sb}_2\text{O}_3$. When heated at $250^\circ \rightarrow \text{Sb}_2\text{Cl}_2\text{O}_5$, and at $>320^\circ \rightarrow \text{Sb}_2\text{O}_3$

References:

Gmelin Handbook Inorg. Chem., Syst. No. 18, 1949, 465, (rev, bibl)

Petrov, K.I. *et al.*, Zh. Neorg. Khim., 1971, **16**, 1895; 1973, **18**, 2922; Russ. J. Inorg. Chem. (Engl. Transl.), 1971, **16**, 1006; 1973, **18**, 1554, (ir, Raman, cryst struct)

Brauer, G., Handbuch Prap. Anorg. Chem., 3rd edn., Ferdinand Enke Verlag, 1975, **1**, 589, (synth)

Glybin, V.P. *et al.*, Zh. Neorg. Khim., 1980, **25**, 313; Russ. J. Inorg. Chem. (Engl. Transl.), 1980, **25**, 170, (struct)