>>**Entry Name: *N*-Benzylhydroxylamine**
Synonym(s): *N*-Hydroxybenzenemethanamine, 9CI

CAS Registry Number: 622-30-0
$PhCH_2NHOH$

Molecular Formula: C_7H_9NO
Molecular Weight: 123.154
Accurate Mass: 123.068414
Percentage Composition: C 68.27%; H 7.37%; N 11.37%; O 12.99%
Use/Importance: Antioxidant. Used in intramolecular nitrone-alkene cycloadditions
Physical Description: Cryst. (petrol)
Melting Point: Mp 57°

Fluka: 13454
Sigma: B2902

>>**Derivative:** Hydrochloride

CAS Registry Number: 29601-98-7
Melting Point: Mp 110°
Aldrich: 36518-1

>>**Derivative:** *N*-Formyl

Molecular Formula: $C_8H_9NO_2$
Molecular Weight: 151.165
Accurate Mass: 151.063329
Percentage Composition: C 63.57%; H 6.00%; N 9.27%; O 21.17%
Physical Description: Needles (Et_2O/petrol)
Melting Point: Mp 49-50°

>>**Derivative:** *N*-Ac

Molecular Formula: $C_9H_{11}NO_2$
Molecular Weight: 165.191
Accurate Mass: 165.078979
Percentage Composition: C 65.44%; H 6.71%; N 8.48%; O 19.37%
Physical Description: Plates
Melting Point: Mp 124°

>>**Derivative:** *N*-Benzoyl

Molecular Formula: $C_{14}H_{13}NO_2$
Molecular Weight: 227.262
Accurate Mass: 227.094629
Percentage Composition: C 73.99%; H 5.77%; N 6.16%; O 14.08%
Physical Description: Needles (Et_2O)

>>**Derivative:** *N*-Nitroso

Molecular Formula: $C_7H_8N_2O_2$
Molecular Weight: 152.152
Accurate Mass: 152.058578
Percentage Composition: C 55.26%; H 5.30%; N 18.41%; O 21.03%
Melting Point: Mp 77-78°

References:

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 564A, *(nmr)*
Feuer, H. *et al.*, *J.A.C.S.*, 1962, **84**, 3771, *(synth)*
Feuer, H. *et al.*, *J.O.C.*, 1965, **30**, 2877, *(synth)*
Isowa, Y. *et al.*, *Bull. Chem. Soc. Jpn.*, 1974, **47**, 720, *(ir, synth)*
Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1981, **9**, 42; 1986, **12**, 13, *(use)*
Maskill, H. *et al.*, *J.A.C.S.*, 1987, **109**, 2062, *(synth, deriv, pmr)*
Kawakani, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 2000, **73**, 2423-2444, *(synth)*