



>>Entry Name: 1,4-Bis(chloromethyl)benzene, 9Cl

Synonym(s): α,α' -Dichloro-*p*-xylene, 8Cl. *p*-Xylylene dichloride

CAS Registry Number: 623-25-6

Molecular Formula: C₈H₈Cl₂

Molecular Weight: 175.057

Accurate Mass: 174.000304

Percentage Composition: C 54.89%; H 4.61%; Cl 40.50%

Use/Importance: Crosslinking agent for polystyrene

Physical Description: Platelets (hexane or EtOH)

Melting Point: Mp 104.1-104.3° (100°)

Boiling Point: Bp 240-245° dec.. Bp₁₂ 135°

Other Data: Undergoes catalytic condensation polym. to give poly(*p*-xylylidene)

Hazard & Toxicity: Eye and skin irritant. LD₅₀ (rat, orl) 1780 mg/kg

Aldrich: 10574-0

Fluka: 36586

References:

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 993A, (*ir*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 911C, (*ir*)

Aldrich Library of NMR Spectra, 2nd edn., 1983, **1**, 798B, (*nmr*)

Quelet, R., *Bull. Soc. Chim. Fr.*, 1933, **53**, 222

Kulka, M., *Can. J. Res., Sect. B*, 1945, **23**, 106, (*synth*)

Farthing, A.C., *J.C.S.*, 1953, 3261

Errede, L.A. *et al.*, *J.A.C.S.*, 1960, **82**, 463

Hoyt, J.M. *et al.*, *Macromol. Synth.*, 1968, **3**, 122, (*polym*)

Anderson, S.N. *et al.*, *J.C.S. Perkin 2*, 1972, 311, (*pmr*)

Tsuchida, E. *et al.*, *Makromol. Chem.*, 1972, **151**, 207, (*synth*)

Yamazaki, Y. *et al.*, *Polym. J. (Tokyo)*, 1983, **15**, 905, (*polym*)

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, DGP600