




>>Entry Name: Bis(glycinato)copper, 11Cl, 10Cl, 9Cl, 8Cl



CAS Registry Number: 13479-54-4

Molecular Formula: C₄H₈CuN₂O₄

Molecular Weight: 211.664

Accurate Mass: 210.988006

Percentage Composition: C 22.70%; H 3.81%; Cu 30.02%; N 13.23%; O 30.24%

Use/Importance: Used in photometric detn. of Cu and in catalysis

>>Variant: *cis*-form

CAS Registry Number: 10171-08-1

Molecular Formula: C₄H₈CuN₂O₄

Molecular Weight: 211.664

Accurate Mass: 210.988006

Percentage Composition: C 22.70%; H 3.81%; Cu 30.02%; N 13.23%; O 30.24%

>>Derivative: Monohydrate

CAS Registry Number: 23678-02-6

Physical Description: Dark blue needles

>>Variant: *trans*-form

CAS Registry Number: 10170-99-7

Molecular Formula: C₄H₈CuN₂O₄

Molecular Weight: 211.664

Accurate Mass: 210.988006

Percentage Composition: C 22.70%; H 3.81%; Cu 30.02%; N 13.23%; O 30.24%

>>Derivative: Monohydrate

CAS Registry Number: 14932-09-3

Physical Description: Deep blue rhombic cryst.

Melting Point: Mp 228 dec.°

Solubility: Sol. H₂O, sl. sol. EtOH

>>Derivative: Dihydrate

CAS Registry Number: 23678-01-5

Physical Description: Light blue powdery cryst.

Melting Point: Mp 225 dec.°

Solubility: Sol. H₂O

References:

Tomita, V.K., *Bull. Chem. Soc. Jpn.*, 1961, **34**, 280, (*cryst struct, synth*)

Herlinger, A.W. *et al.*, *J.A.C.S.*, 1970, **92**, 6474, (*ir*)

Allen, H.C. *et al.*, *J. Chem. Phys.*, 1972, **56**, 997, (*epr*)

Hayakawa, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1973, **46**, 2792, (*use*)

Rupp, H. *et al.*, *Bioinorg. Chem.*, 1976, **6**, 45, (*pe*)

O'Brien, P., *J. Chem. Educ.*, 1982, **59**, 1052, (*synth, ir*)

Merck Index, 11th edn., 1989, No. 2644