




>>Entry Name: Bis(*N,N*-dibutyldithiocarbamato)nickel(II)

Synonym(s): Bis(dibutylcarbamodithioato-*S,S'*)nickel, 11Cl



CAS Registry Number: 13927-77-0

Molecular Formula: C₁₈H₃₆N₂NiS₄

Molecular Weight: 467.45

Accurate Mass: 466.111474

Percentage Composition: C 46.25%; H 7.76%; N 5.99%; Ni 12.56%; S 27.44%

General Statement: Square planar monomer; constrast struct. of Bis(*N,N*-dibutylthiocarbamato)nickel(II) [JQV48-V](#)

Use/Importance: Antioxidant and antiozonant for dyes

Physical Description: Green cryst. (C₆H₆/EtOH)

Melting Point: Mp 91°

Solubility: Sol. C₆H₆, Me₂CO

Other Data: Diamagnetic

References:

- Cavell, H.J. *et al.*, *J.C.S.*, 1935, 621, (*synth*)
Malatesta, L., *Gazz. Chim. Ital.*, 1937, **67**, 738, (*synth, magnetism*)
Talaridis, A. *et al.*, *Fresenius' Z. Anal. Chem.*, 1978, **293**, 211, (*synth, gc*)
Oktavec, D. *et al.*, *Coll. Czech. Chem. Comm.*, 1980, **45**, 1495, (*ir*)
Riekkola, M.L., *Finn. Chem. Lett.*, 1980, 83, (*gc*)
Riekkola, M.L., *Acta Chem. Scand., Ser. A*, 1983, **37**, 691, (*ms*)
Lokaj, J. *et al.*, *Chem. Zvesti*, 1984, **38**, 313, (*cryst struct*)
Payne, R. *et al.*, *J. Electron Spectrosc. Relat. Phenom.*, 1985, **35**, 113, (*pe, ir*)
Matsui, M. *et al.*, *J. Soc. Dyers Colour.*, 1985, **101**, 334, (*use*)
Tersumi, T. *et al.*, *Polyhedron*, 1986, **5**, 707, (*synth*)
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, BIW750