>>**Entry Name: Binedaline,** INN
Synonym(s): *N,N,N'*-Trimethyl-*N'*-(3-phenyl-1*H*-indol-1-yl)-1,2-ethanediamine, 9CI. 1-[[2-(Dimethylamino)ethyl]methylamino]-3-phenylindole. Binodaline. Scha 1659. Sgd 20578. Sgd Scha 1059



CAS Registry Number: 60662-16-0

Molecular Formula: $C_{19}H_{23}N_3$
Molecular Weight: 293.411
Accurate Mass: 293.189197
Percentage Composition: C 77.78%; H 7.90%; N 14.32%
Use/Importance: Noradrenaline uptake inhibitor. Antidepressant agent
Development Status: Never marketed
Hazard & Toxicity: LD_{50} (rat, orl) 1160 mg/kg

>>**Derivative:** Hydrochloride
Synonym(s): Ixprim

CAS Registry Number: 57647-35-5
Melting Point: Mp 188-190°
Hazard & Toxicity: LD_{50} (rat, orl) 1160 mg/kg. LD_{50} (rat, ivn) 26 mg/kg

References:

Ger. Pat., 1975, *Siegfried*, 2 512 702; *CA*, **84**, 30886v, *(synth)*
Schatz, F. *et al.*, *Arzneim.-Forsch.*, 1980, **30**, 919, *(synth, pharmacol)*
Jahn, U. *et al.*, *Arzneim.-Forsch.*, 1983, **33**, 726, *(pharmacol, tox)*
Morin, D. *et al.*, *J. Pharmacol. Exp. Ther.*, 1989, **249**, 288, *(pharmacol)*
Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 250
Lewis, R.J.*, Sax's Dangerous Properties of Industrial Materials, 8th edn., Van Nostrand Reinhold*, 1992, BGC500

https://www.baizuoyan.com - CopyRight ©baizuoyan