




>>Entry Name: Bis(2-aminophenyl) disulfide

Synonym(s): 2,2'-Dithiobisbenzenamine, 9Cl. 2,2'-Dithiodianiline, 8Cl. 2,2'-Diaminodiphenyl disulfide. Intramine



CAS Registry Number: 1141-88-4

Molecular Formula: C₁₂H₁₂N₂S₂

Molecular Weight: 248.372

Accurate Mass: 248.044188

Percentage Composition: C 58.03%; H 4.87%; N 11.28%; S 25.82%

Use/Importance: Vulcanizing agent for urethane rubbers, shows bactericidal and fungicidal activity. Used as EtOH soln. for photometric detn. of Pt(IV) ($\lambda_{\max}$ 725 nm, ϵ 54800)

Physical Description: Leaflets or needles (EtOH aq.)

Melting Point: Mp 93°

Solubility: Sol. acids, EtOH; spar. sol. H₂O

Hazard & Toxicity: Severe eye irritant. LD₅₀ (rat, orl) 558 mg/kg

Aldrich: 16676-6

Fluka: 43767

>>Derivative: Dipicrate

Melting Point: Mp 141°

>>Derivative: 2,2'-N-Di-Ac

CAS Registry Number: 3095-79-2

Molecular Formula: C₁₆H₁₆N₂O₂S₂

Molecular Weight: 332.447

Accurate Mass: 332.065318

Percentage Composition: C 57.81%; H 4.85%; N 8.43%; O 9.63%; S 19.29%

Melting Point: Mp 156°

References:

- Aldrich Library of FT-IR Spectra, 1st edn.*, 1985, **1**, 1197A, (*ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 465B, (*nmr*)
Gialdi, F. *et al.*, *Farmaco, Ed. Sci.*, 1959, **14**, 216, (*synth*)
Davara, A.C., *CA*, 1968, **68**, 95438, (*synth, nmr, ir*)
Uemura, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1977, **50**, 220, (*synth*)
Shapiro, B.L. *et al.*, *J. Phys. Chem. Ref. Data*, 1977, **6**, 919, (*nmr*)
Bag, S.P. *et al.*, *Talanta*, 1977, **24**, 128, (*use*)
Higashi, L.S. *et al.*, *J.A.C.S.*, 1978, **100**, 8101, (*cryst struct*)
Kaitner, B. *et al.*, *Acta Cryst. C*, 1997, **53**, 1103-1105, (*cryst struct*)
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, DXJ800