




>>Entry Name: Acetylferrocene, 9CI

Synonym(s): Ferrocenyl methyl ketone



CAS Registry Number: 1271-55-2

Related CAS Registry Numbers(s): 12193-78-1

Molecular Formula: $C_{12}H_{12}FeO$

Molecular Weight: 228.073

Accurate Mass: 228.023754

Percentage Composition: C 63.20%; H 5.30%; Fe 24.49%; O 7.02%

Source/Synthesis: Synth. from ferrocene and acetic anhydride with various catalysts

Use/Importance: Curing agent in resin production; reduces flammability and suppresses smoke production in pvc

Physical Description: Orange cryst. (heptane)

Melting Point: Mp 85-86°

pKa Value: $pK_{a1} -2.8$ (H_2SO_4 aq.)

NMR Spectra: Pmr 2.28 (CH_3) ppm

Other Data: Exhibits phase transitions at 80° and 3°

Hazard & Toxicity: LD₅₀ (oral, female rats) <5 mg/kg, (oral, male rats) <50 mg/kg, (oral, monkeys) <100 mg/kg

Aldrich: 10686-0

Fluka: 1130

Sigma: A1011

>>Derivative: (E)-Oxime

CAS Registry Number: 32662-56-9

Molecular Formula: $C_{12}H_{13}FeNO$

Molecular Weight: 243.088

Accurate Mass: 243.034653

Percentage Composition: C 59.29%; H 5.39%; Fe 22.97%; N 5.76%; O 6.58%

Physical Description: Orange-yellow flakes (hexane)

Melting Point: Mp 174-175°

>>Derivative: (Z)-Oxime

CAS Registry Number: 32662-57-0

Physical Description: Orange-yellow flakes (hexane)

Melting Point: Mp 144-145°

References:

- Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **2**, 1151B, (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 770B, (nmr)
Rinehart, K.L. *et al.*, *J.A.C.S.*, 1957, **79**, 2749, (ir)
Graham, P.J. *et al.*, *J.A.C.S.*, 1957, **79**, 3416, (synth)
Broadhead, G.D. *et al.*, *J.C.S.*, 1958, 650, (synth)
Lundquist, R.T. *et al.*, *J.O.C.*, 1962, **27**, 1167, (uv)
Org. React. (N.Y.), 1969, **17**, 1, (synth)
Yamakawa, K. *et al.*, *Tetrahedron*, 1970, **26**, 4483, (oxime, synth, ir, pmr)
Olah, G.A. *et al.*, *J. Organomet. Chem.*, 1973, **60**, 511, (pmr)
Braun, S. *et al.*, *J. Organomet. Chem.*, 1975, **97**, 429, (cmr)
Sato, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1984, **57**, 2361, (cryst struct, Mössbauer)
U.S. Pat., 1985, 4 525 232; *CA*, 1985, **103**, 142954h, (use)
Leung, H.W. *et al.*, *Toxicol. Lett.*, 1987, **38**, 103, (tox)
Carty, P. *et al.*, *Appl. Organomet. Chem.*, 1990, **4**, 127, (use)
Houlton, A. *et al.*, *J.C.S. Dalton*, 1991, 467, (cmr, Mössbauer, Fe-57 nmr)
Synthetic Methods of Organometallic and Inorganic Chemistry, (ed. Hermann, W.A.), Thieme, 1996, **1**, 136-137, (synth)
Darin, V.A. *et al.*, *J. Prakt. Chem.*, 1999, **341**, 588-591, (synth, cmr)
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ABA750