



Percentage Composition: C 42.86%; H 4.32%; N 29.99%; O 22.84%

Physical Description: Cryst. (EtOH)

Melting Point: Mp 245-247°

Optical Rotation: $[\alpha]_D^{25} -29$ (c, 1.08 in 1M HCl)

>>**Derivative:** *N*-Methylamide

CAS Registry Number: 35788-27-3

Molecular Formula: C₁₁H₁₄N₆O₄

Molecular Weight: 294.269

Accurate Mass: 294.107654

Percentage Composition: C 44.90%; H 4.80%; N 28.56%; O 21.75%

Physical Description: Cryst. (MeOH)

Melting Point: Mp 241-243°

>>**Derivative:** *N*-Ethylamide

Synonym(s): 1-(6-Amino-9*H*-purin-9-yl)-1-deoxy-*N*-ethyl-β-D-ribofuranuronamide, 9Cl. Adenosine-5'-*N*-ethyluronamide. 5'-*N*-Ethylcarboxamideadenosine. NECA

CAS Registry Number: 35920-39-9

Molecular Formula: C₁₂H₁₆N₆O₄

Molecular Weight: 308.296

Accurate Mass: 308.123304

Percentage Composition: C 46.75%; H 5.23%; N 27.26%; O 20.76%

Biological Use/Importance: Non-selective adenosine receptor agonist

Physical Description: Cryst. (EtOH or H₂O)

Melting Point: Mp 249-250° (245-246°)

Optical Rotation: $[\alpha]_D^{26} -16.3$ (c, 0.92 in HCl)

>>**Derivative:** *N*-Ethylamide, 2,3-di-Ac

CAS Registry Number: 58048-26-3

Molecular Formula: C₁₆H₂₀N₆O₆

Molecular Weight: 392.371

Accurate Mass: 392.144434

Percentage Composition: C 48.98%; H 5.14%; N 21.42%; O 24.47%

Melting Point: Mp 96-102°

Optical Rotation: $[\alpha]_D^{26} -19$ (c, 1.8 in H₂O)

>>**Derivative:** *N,N*-Diethylamide

Molecular Formula: C₁₄H₂₀N₆O₄

Molecular Weight: 336.35

Accurate Mass: 336.154604

Percentage Composition: C 49.99%; H 5.99%; N 24.99%; O 19.03%

Physical Description: Cryst. (MeOH)

Melting Point: Mp 244-245°

References:

- Stein, H.H., *J. Med. Chem.*, 1973, **16**, 1306, (*synth*)
Meris, I. *et al.*, *Xenobiotica*, 1973, **3**, 381, (*metab*)
U.S. Pat., 1974, *Abbott*, 3 855 205; *CA*, **82**, 171372v, (*synth*)
Stein, H.H. *et al.*, *Ann. N.Y. Acad. Sci.*, 1975, **255**, 380, (*rev, pharmacol*)
Prasad, R.N. *et al.*, *J. Med. Chem.*, 1976, **19**, 1180; 1980, **23**, 313-319, (*synth, derivs, bibl*)
Raberger, G. *et al.*, *Arch. Int. Pharmacodyn. Ther.*, 1977, **230**, 140-149, (*NECA, pharmacol*)
Prasad, R.N. *et al.*, *J. Med. Chem.*, 1980, **23**, 313-319, (*NECA, synth, pharmacol*)
Olsson, R.A. *et al.*, *J. Med. Chem.*, 1986, **29**, 1683-1689, (*synth*)
Moos, W.H. *et al.*, *Nucleosides Nucleotides*, 1989, **8**, 449-461, (*cryst struct, ethylamide*)
Bridges, A.J. *et al.*, *Nucleic Acid Chem.*, 1991, **4**, 230-239, (*NECA, synth, derivs, pharmacol*)
Wnuk, S.F. *et al.*, *J. Med. Chem.*, 1996, **39**, 4162-4166, (*benzyl ester, amide derivs, synth, pmr*)
Sreekante, G. *et al.*, *Acta Cryst. C*, 1998, **54**, 647-649, (*cryst struct*)