



CAS Registry Number: 13707-88-5

Melting Point: Mp 108-110°

Hazard & Toxicity: LD₅₀ (rat, orl) 590 mg/kg

Sigma: A8676

References:

- Brandstrom, A. *et al.*, *Acta Pharm. Suec.*, 1966, **3**, 303-310; *CA*, **68**, 21622, (*synth*)
Karow, M.A. *et al.*, *Prog. Drug Res.*, 1971, **15**, 103, (*rev*)
Barrans, Y. *et al.*, *Acta Cryst. B*, 1973, **29**, 1264, (*cryst struct*)
Dangoumau, J. *et al.*, *J. Pharmacol.*, 1973, **4**, 5, (*conformn, props*)
Caron, M.G. *et al.*, *J. Biol. Chem.*, 1976, **25**, 2374-2384, (*synth, dihydro*)
Randall, M.H. *et al.*, *J. Med. Chem.*, 1977, **20**, 1090-1094, (*S-form dihydro*)
Aablad, B. *et al.*, *Pharmacol. Antihypertens. Drugs*, 1980, 223, (*rev, pharmacol*)
Negwer, M., *Organic-Chemical Drugs and their Synonyms*, 6th edn., *Akademie-Verlag*, 1987, 3726, (*synonyms*)
Landi, M. *et al.*, *Biochem. Pharmacol.*, 1992, **44**, 665-672, (*bibl, dihydro*)
Martindale, The Extra Pharmacopoeia, 30th edn., *Pharmaceutical Press*, 1993, 628
Höld, K.M., *J. Chromatogr. Sci.*, 1996, **34**, 13-19, (*glc, ms*)
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., *Van Nostrand Reinhold*, 1992, AGW000; AGW250