



>>>Derivative: *N,N*-Di-Et

Synonym(s): *N,N*-Diethyl-4-[(4-nitrophenyl)azo]benzenamine, 9Cl

CAS Registry Number: 3025-52-3

Extra CAS Registry Numbers(s): 80631-72-7 80631-73-8

Molecular Formula: C₁₆H₁₈N₄O₂

Molecular Weight: 298.344

Accurate Mass: 298.142976

Percentage Composition: C 64.41%; H 6.08%; N 18.78%; O 10.73%

Physical Description: Dark steel blue needles (C₆H₆); red brown plates (EtOH)

Melting Point: Mp 150-152° (151°)

>>>Derivative: *N*-Ph

Synonym(s): 4-[(4-Nitrophenyl)azo]-*N*-phenylbenzenamine, 9Cl. C.I. Disperse Orange 1. C.I. 11080

CAS Registry Number: 2581-69-3

Extra CAS Registry Numbers(s): 82248-50-8

Molecular Formula: C₁₈H₁₄N₄O₂

Molecular Weight: 318.334

Accurate Mass: 318.111676

Percentage Composition: C 67.92%; H 4.43%; N 17.60%; O 10.05%

Physical Description: Dark red cryst (C₆H₆/hexane or EtOH aq.)

Melting Point: Mp 162.2-163° (152-153°)

>>>Derivative: *N*-(2-Hydroxyethyl), *N*-Et

Synonym(s): 2-[Ethyl[4-[(4-nitrophenyl)azo]phenyl]amino]ethanol, 9Cl. 4-[Ethyl(2-hydroxyethyl)amino]-4'-nitroazobenzene. Disperse Fast Scarlet B. C.I. Disperse Red 1. C.I. Solvent Red 14

CAS Registry Number: 2872-52-8

Molecular Formula: C₁₆H₁₈N₄O₃

Molecular Weight: 314.343

Accurate Mass: 314.137891

Percentage Composition: C 61.14%; H 5.77%; N 17.82%; O 15.27%

Use/Importance: Dye for polyesters, also used in manuf. of nonlinear optical polymers

Physical Description: Scarlet platelets (Me₂CO)

Melting Point: Mp 163°

References:

- Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **2**, 966C, (*ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 547C, (*nmr*)
Meldola, R., *J.C.S.*, 1883, **43**, 425-442; 1884, **45**, 106-121, (*N,N*-di-Me, *N*-Ph)
Bamberger, E., *Ber.*, 1895, **28**, 837-854, (*N,N*-di-Me, *N,N*-di-Et)
Witt, O.N. *et al.*, *Ber.*, 1912, **45**, 1148, (*synth*)
Meyer, K.H. *et al.*, *Ber.*, 1921, **54**, 2272, (*synth*)
Pongratz, A. *et al.*, *Ber.*, 1938, **71**, 1287-1296, (*N,N*-di-Me, *synth*, *uv*)
v. Mechel, L. *et al.*, *Helv. Chim. Acta*, 1941, **24**, 151E, (*synth*)
Knight, A.H. *et al.*, *J. Soc. Dyers Colour.*, 1950, **66**, 169-177, (*manuf*, *deriv*)
Wahl, H. *et al.*, *Bull. Soc. Chim. Fr.*, 1952, 366-369, (*derivs*, *synth*)
Le Fevre, R.J.W. *et al.*, *Aust. J. Chem.*, 1957, **10**, 26-33, (*N,N*-di-Me, *ir*)
Marple, L.W. *et al.*, *Anal. Chem.*, 1963, **35**, 1305, (*use*)
Fritz, J.S. *et al.*, *Talanta*, 1966, **13**, 939, (*use*)
Gilland, J.C. *et al.*, *Org. Mass Spectrom.*, 1974, **9**, 1148-1153, (*ms*, *derivs*)
Rondestvedt, C.S., *J.O.C.*, 1977, **42**, 1786-1790, (*N*-Ph)
Asano, T. *et al.*, *J.A.C.S.*, 1981, **103**, 5161-5165; 1982, **104**, 4900-4904, (*N,N*-di-Me, *N*-Ph)
Nishimura, N. *et al.*, *J.C.S. Perkin 2*, 1986, 1839-1845, (*N,N*-di-Et, *E/Z isom*)
Savarino, P. *et al.*, *Dyes Pigm.*, 1990, **13**, 71-80, (*cmr*, *derivs*)
Sigma-Aldrich Library of Stains, Dyes and Indicators, 287