



Percentage Composition: C 71.46%; H 8.99%; N 5.95%; O 13.60%

Use/Importance: Local anaesthetic

Calculated Log P Data: Log P 3.37 (calc)

>>**Derivative:** *N*-Propyl, *O*-benzoyl; hydrochloride

Synonym(s): Meprylcaine hydrochloride, USAN. Oracaine. Marvacaine. Meprocaïne

CAS Registry Number: 956-03-6

Physical Description: Cryst. (2-propanol)

Melting Point: Mp 150-152°

Hazard & Toxicity: LD₅₀ (mus, scu) 262 mg/kg

Sigma: M4281

>>**Derivative:** *N*-Benzyl

CAS Registry Number: 10250-27-8

Molecular Formula: C₁₁H₁₇NO

Molecular Weight: 179.261

Accurate Mass: 179.131014

Percentage Composition: C 73.70%; H 9.56%; N 7.81%; O 8.93%

Physical Description: Cryst. (cyclohexane)

Melting Point: Mp 54-56°

References:

- Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **1**, 539B, (nmr)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 425D, (ir)
Aldrich Library of Infrared Spectra, 3rd edn., 1981, 198C, (ir)
Johnson, K. *et al.*, *J.O.C.*, 1943, **8**, 7, (synth)
U.S. Pat., 1946, 2 404 319; *CA*, **40**, 62144, (Bufylline)
U.S. Pat., 1956, 2 767 207; *CA*, **51**, 583f, (Meprylcaine)
Thompson, W.E. *et al.*, *J. Pharm. Sci.*, 1965, **54**, 1819, (ir, derivs)
Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1967, **1**, 37; 1972, **3**, 14; 1977, **6**, 20; 368; 1974, **4**, 540, (use)
Leonard, N.J. *et al.*, *J.O.C.*, 1967, **32**, 3607, (N,N-di-Me)
Lichter, R.L. *et al.*, *J.A.C.S.*, 1972, **94**, 2495, (nmr)
Sameski, J.E. *et al.*, *Anal. Chem.*, 1975, **47**, 2116, (cmr)
Kirk-Othmer Encycl. Chem. Technol., 4th edn., Wiley, 1991, **2**, 26, (rev, use, tox)
Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 1015
Urleb, U. *et al.*, *Synth. Commun.*, 1994, **24**, 1415, (N-benzyl, synth)
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, IIA000; OJI600