




>>Entry Name: Di(2-butyl)amine

Synonym(s): *N*-(1-Methylpropyl)-2-butanamine, 9Cl. Di-*sec*-butylamine, 8Cl. 1,1'-Dimethyldipropylamine. 2,2'-Iminobisbutane



CAS Registry Number: 626-23-3

Molecular Formula: C₈H₁₉N

Molecular Weight: 129.245

Accurate Mass: 129.151749

Percentage Composition: C 74.35%; H 14.82%; N 10.84%

General Statement: Long known as diastereoisomeric mixtures. Prominence given here to the most recent preparation of pure diastereoisomers

Physical Description: Liq.

Boiling Point: Bp 132-134°

Solubility: V. sol. H₂O

Density: d₄⁰ 0.783

Hazard & Toxicity: Flammable, fl. p. 24°. Irritant

>>Variant: (S,S)-form

CAS Registry Number: 68852-54-0

Molecular Formula: C₈H₁₉N

Molecular Weight: 129.245

Accurate Mass: 129.151749

Percentage Composition: C 74.35%; H 14.82%; N 10.84%

>>Derivative: Hydrochloride

CAS Registry Number: 131491-40-2

Melting Point: Mp 176°

Optical Rotation: [α]_D²³ +3.42 (c, 10 in MeOH) (>=99% ee)

>>Variant: (RS,SR)-form

Synonym(s): *meso*-form

CAS Registry Number: 39056-20-7

Molecular Formula: C₈H₁₉N

Molecular Weight: 129.245

Accurate Mass: 129.151749

Percentage Composition: C 74.35%; H 14.82%; N 10.84%

>>Derivative: Hydrochloride

CAS Registry Number: 131565-41-8

Melting Point: Mp 122°

References:

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 476C, (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 381A, (*ir*)

Stewart, J.E., *J. Chem. Phys.*, 1959, **30**, 1259, (*ir*)

Gohlke, R.S. *et al.*, *Anal. Chem.*, 1962, **34**, 1281, (*ms*)

Brown, H.C. *et al.*, *J.O.C.*, 1991, **56**, 1170, (*synth, pmr, cmr*)

Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DDT600

Bretherick, L., *Handbook of Reactive Chemical Hazards*, 4th edn., Butterworths, 1990, 2887