



Molecular Weight: 173.301
Accurate Mass: 173.189197
Percentage Composition: C 62.38%; H 13.38%; N 24.25%
Physical Description: No phys. props. reported

>>**Derivative:** $N^3, N^3, N^{3'}, N^{3'}$ -Tetra-Me

Synonym(s): N' -[3-(Dimethylamino)propyl]- N, N -dimethyl-1,3-propanediamine. 3,3'-Bis(dimethylamino)dipropylamine

CAS Registry Number: 6711-48-4
Molecular Formula: $C_{10}H_{25}N_3$
Molecular Weight: 187.328
Accurate Mass: 187.204847
Percentage Composition: C 64.12%; H 13.45%; N 22.43%
Melting Point: Mp -78°
Boiling Point: Bp₂₀ 128-131°
Refractive Index: n_D^{20} 1.4490

Aldrich: 34855-4

>>**Derivative:** N -Penta-Me

Synonym(s): N -[3-(Dimethylamino)propyl]- N, N', N' -trimethyl-1,3-propanediamine, 9Cl. 3,3'-Bis(dimethylamino)- N -methyldipropylamine, 8Cl. N, N, N', N'', N'' -Pentamethyldipropylenetriamine

CAS Registry Number: 3855-32-1
Molecular Formula: $C_{11}H_{27}N_3$
Molecular Weight: 201.354
Accurate Mass: 201.220497
Percentage Composition: C 65.62%; H 13.52%; N 20.87%
Use/Importance: Catalyst for polyurethane foam production; auxiliary for lithiation reactions
Physical Description: Liq.
Boiling Point: Bp₁₁ 100-102°

References:

- Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **1**, 309C; 310B, (*ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 500B; 500C; 501A, (*nmr*)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 397A; 397B, (*ir*)
Wiedeman, D.F. *et al.*, *J.A.C.S.*, 1945, **67**, 1994, (*synth*)
Terent'ev, A.P. *et al.*, *Zh. Obshch. Khim.*, 1950, **20**, 1073; *CA*, **44**, 9350a, (*synth*)
Marxer, A. *et al.*, *Helv. Chim. Acta*, 1951, **34**, 924-931, (*N-penta-Me*)
Israel, M. *et al.*, *J. Med. Chem.*, 1964, **7**, 710, (*synth*)
Schrier, M. *et al.*, *Mikrochim. Acta*, 1965, 1091; 1967, 218, (*use*)
Ger. Pat., 1978, 2 739 353; *CA*, **89**, 111389r, (*N-penta-Me*)
Bergeron, R.J. *et al.*, *Synthesis*, 1981, 732, (*synth*)
Canadian Pat., 1982, 1 132 292; *CA*, **98**, 55043v, (*N-penta-Me*)
Murahashi, S. *et al.*, *J.A.C.S.*, 1983, **105**, 5002, (*synth*, *pmr*, *ir*)
Dagnall, S.P. *et al.*, *J.C.S. Perkin 2*, 1984, 435, (*cmr*)
Hamana, K. *et al.*, *J. Biochem. (Tokyo)*, 1985, **97**, 1595; 1991, **109**, 444, (*occur*)
Tanaka, Y. *et al.*, *CA*, 1988, **109**, 232044q, (*crosslinking*)
Carboni, B. *et al.*, *Tet. Lett.*, 1988, **29**, 1279, (*synth*)
Yamamoto, S. *et al.*, *J. Biochem. (Tokyo)*, 1993, **113**, 538, (*bisdihydroxybenzoyl*)
Thurner, A. *et al.*, *Synth. Commun.*, 1998, **28**, 443-449, (*N-penta-Me*, *use*)
Lewis, R.J., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AIX250; BGU750; TDP750